



www.theoryofgroups.ir

---

**International Journal of Group Theory**  
ISSN (print): 2251-7650, ISSN (on-line): 2251-7669  
Vol. 8 No. 1 (2019), pp. 35-42.  
© 2019 University of Isfahan

---



www.ui.ac.ir

## FINITE GROUPS OF THE SAME TYPE AS SUZUKI GROUPS

SEYED HASSAN ALAVI, ASHRAF DANESHKHAH\* AND HOSEIN PARVIZI MOSAED

Communicated by Alireza Abdollahi

**ABSTRACT.** For a finite group  $G$  and a positive integer  $n$ , let  $G(n)$  be the set of all elements in  $G$  such that  $x^n = 1$ . The groups  $G$  and  $H$  are said to be of the same (order) type if  $|G(n)| = |H(n)|$ , for all  $n$ . The main aim of this paper is to show that if  $G$  is a finite group of the same type as Suzuki groups  $Sz(q)$ , where  $q = 2^{2m+1} \geq 8$ , then  $G$  is isomorphic to  $Sz(q)$ . This addresses to the well-known J. G. Thompson's problem (1987) for simple groups.

### 1. Introduction

For a finite group  $G$  and a positive integer  $n$ , let  $G(n)$  consist of all elements  $x$  satisfying  $x^n = 1$ . The *order type* of  $G$  is defined to be the function whose value at  $n$  is the order of  $G(n)$ . In 1987, J. G. Thompson [11, Problem 12.37] posed a problem which is related to algebraic number fields:

Is it true that a group is solvable if its type is the same as that of a solvable one?

This problem links to the set  $nse(G)$  of *the number of elements of the same order* in  $G$ . Indeed, it turns out that if two groups  $G$  and  $H$  are of the same type, then  $nse(G) = nse(H)$  and  $|G| = |H|$ . Therefore, if a group  $G$  has been uniquely determined by its order and  $nse(G)$ , then Thompson's problem is true. One may ask this problem for non-solvable groups, in particular, finite simple groups. In this direction, Shao et al [13] studied finite simple groups whose order is divisible by at most four primes. Following this investigation, such problem has been studied for some families

---

MSC(2010): Primary: 20D06; Secondary: 20E32, 20E34.

Keywords: Suzuki group, Thompson's problem, element order.

Received: 08 March 2017, Accepted: 14 July 2017.

\*Corresponding author.

<http://dx.doi.org/10.22108/ijgt.2017.21556>

of simple groups [1, 2, 3, 13] including Suzuki groups  $Sz(q)$  with  $q$  prime [10]. In this paper, we generalize the main result in [10] and prove that

**Theorem 1.1.** *Let  $G$  be a group with  $nse(G) = nse(Sz(q))$  and  $|G| = |Sz(q)|$ . Then  $G$  is isomorphic to  $Sz(q)$ .*

As noted above, as an immediate consequence of Theorem 1.1, we have that

**Corollary 1.2.** *If  $G$  is a finite group of the same type as  $Sz(q)$ , then  $G$  is isomorphic to  $Sz(q)$ .*

In order to prove Theorem 1.1, we use a partition of Suzuki groups  $S := Sz(q)$ , where  $q = 2^{2m+1} \geq 8$ , see Lemma 3.2, that is to say, a set of subgroups  $H_i$  of  $S$ , for  $i = 1, \dots, s$ , such that each nontrivial element of  $S$  belongs to exactly one subgroup  $H_i$ . We use this information to determine the set  $nse(S)$  in Proposition 3.3. It is also a main tool to show that 2 is an isolated vertex in the prime graph of a group  $G$  satisfying hypotheses of Theorem 1.1, see Proposition 4.1. Then we show that  $G$  is neither Frobenius, nor 2-Frobenius group. Finally, we obtain a section of  $G$  which is isomorphic to  $S$  and prove that  $G$  is isomorphic to  $S$ .

Finally, we give some brief comments on the notation used in this paper. Throughout this article all groups are finite. Our group-theoretic notation is standard, and it is consistent with the notation in [4, 6, 7]. We denote a Sylow  $p$ -subgroup of  $G$  by  $G_p$ . We also use  $n_p(G)$  to denote the number of Sylow  $p$ -subgroups of  $G$ . For a positive integer  $n$ , the set of prime divisors of  $n$  is denoted by  $\pi(n)$ , and if  $G$  is a finite group,  $\pi(G) := \pi(|G|)$ , where  $|G|$  is the order of  $G$ . We denote the set of elements' orders of  $G$  by  $\omega(G)$  known as *spectrum* of  $G$ . The *prime graph*  $\Gamma(G)$  of a finite group  $G$  is a graph whose vertex set is  $\pi(G)$ , and two vertices  $p$  and  $q$  are adjacent if and only if  $pq \in \omega(G)$ . Assume further that  $\Gamma(G)$  has  $t(G)$  connected components  $\pi_i$ , for  $i = 1, 2, \dots, t(G)$ . The positive integers  $n_i$  with  $\pi(n_i) = \pi_i$  are called order components of  $G$ . Clearly,  $|G| = n_1 \cdots n_{t(G)}$ . In the case where  $G$  is of even order, we always assume that  $2 \in \pi_1$ , and  $\pi_1$  is said to be the even component of  $G$ . In this way,  $\pi_i$  and  $n_i$  are called odd components and odd order components of  $G$ , respectively. Recall that  $nse(G)$  is the set of the number of elements in  $G$  with the same order. In other word,  $nse(G)$  consists of the numbers  $m_i(G)$  of elements of order  $i$  in  $G$ , for  $i \in \omega(G)$ . Here,  $\phi$  is the *Euler totient* function.

## 2. Preliminaries

In this section, we introduce the some known results which will be used in the proof of the main result.

**Lemma 2.1.** [8, Theorem 9.1.2] *Let  $G$  be a finite group, and let  $n$  be a positive integer dividing  $|G|$ . Then  $n$  divides  $|G(n)|$ .*

The proof of the following result is straightforward by Lemma 2.1. Recall that  $nse(G) = \{m_i(G) \mid i \in \omega(G)\}$ .

**Lemma 2.2.** *Let  $G$  be a finite group. Then for every  $i \in \omega(G)$ ,  $\phi(i)$  divides  $m_i(G)$ , and  $i$  divides  $\sum_{j|i} m_j(G)$ . Moreover, if  $i > 2$ , then  $m_i(G)$  is even.*

**Lemma 2.3** (Theorem 3 in [15]). *Let  $G$  be a finite group of order  $n$ . Then the number of elements whose orders are multiples of  $t$  is either zero, or a multiple of the greatest divisor of  $n$  that is prime to  $t$ .*

In what follows, recall that  $t(G)$  is the number of connected components of the prime graph  $\Gamma(G)$ .

**Lemma 2.4.** [5, Theorem 1] *Let  $G$  be a Frobenius group of even order with kernel  $K$  and complement  $H$ . Then  $t(G) = 2$ ,  $\pi(H)$  and  $\pi(K)$  are vertex sets of the connected components of  $\Gamma(G)$ .*

A group  $G$  is called 2-Frobenius if there exists a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $G/H$  and  $K$  are Frobenius groups with kernel  $K/H$  and  $H$  respectively.

**Lemma 2.5.** [5, Theorem 2] *Let  $G$  be a 2-Frobenius group of even order. Then  $t(G) = 2$ ,  $\pi(G/K) \cup \pi(H) = \pi_1$ ,  $\pi(K/H) = \pi_2$ , and  $G/K$  and  $K/H$  are cyclic groups and  $|G/K|$  divides  $|\text{Aut}(K/H)|$ .*

### 3. Elements of the same order in Suzuki groups

In this section, we determine the set of the number of elements of the same order in Suzuki groups.

**Lemma 3.1** ([14]). *Let  $S = \text{Sz}(q)$  with  $q = 2^{2m+1} \geq 8$ . Then  $\omega(S)$  consists of all factors of  $4$ ,  $q - 1$  and  $q \pm \sqrt{2q} + 1$ .*

Let  $G$  be a group, and let  $H_1, \dots, H_t$  be subgroups of  $G$ . Then the set  $\{H_1, \dots, H_t\}$  forms a partition of  $G$  if each non-trivial element of  $G$  belongs to exactly one subgroup  $H_i$  of  $G$ . Lemma 3.2 below introduces a partition of Suzuki groups.

**Lemma 3.2.** *Let  $S = \text{Sz}(q)$  with  $q = 2^{2m+1} \geq 8$ , and let  $\mathbb{F} := \text{GF}(q)$ . Then*

- (a)  *$S$  possesses cyclic subgroups  $U_1$  and  $U_2$  of orders  $q + \sqrt{2q} + 1$  and  $q - \sqrt{2q} + 1$ , respectively;*
- (b) *if  $1 \neq u \in U_i$ , for  $i = 1, 2$ , then  $S(u) = U_i$ . Moreover,  $|\mathbf{N}_S(U_i) : U_i| = 4$ ;*
- (c)  *$S$  possesses a cyclic subgroup  $V$  of order  $q - 1$  and  $|\mathbf{N}_S(V) : V| = 2$ ;*
- (d)  *$S$  possesses a 2-subgroup  $W$  of orders  $q^2$  and exponent 4 and  $|S : \mathbf{N}_S(W)| = q^2 + 1$ . Moreover, the elements of  $W$  are of the form*

$$(3.1) \quad w(a, b) := \begin{pmatrix} 1 & 0 & 0 & 0 \\ a & 1 & 0 & 0 \\ b & a\pi & 1 & 0 \\ a^2(a\pi) + ab + b\pi & a(a\pi) + b & a & 1 \end{pmatrix},$$

where  $a, b \in \mathbb{F}$  and  $\pi \in \text{Aut}(\mathbb{F})$  maps  $x$  to  $x^{2m+1}$ , for all  $x \in \mathbb{F}$ .

- (e) *the conjugates of  $U_1, U_2, V$  and  $W$  form a partition of  $S$ .*

*Proof.* All parts of this result follow from Lemma 3.1 and Theorem 3.10 in [9] except for the facts that  $|N_S(V) : V| = 2$  and  $|S : N_S(W)| = q^2 + 1$  which can be found in the proof of Theorem 3.10 in [9].  $\square$

**Proposition 3.3.** *Let  $S = \text{Sz}(q)$  with  $q = 2^{2m+1} \geq 8$ . Then the set  $\text{nse}(S)$  consists of exactly one the following numbers*

- (a)  $1, (q-1)(q^2+1), q(q-1)(q^2+1)$ ;
- (b)  $\phi(i)q^2(q \mp \sqrt{2q} + 1)(q-1)/4$ , where  $i > 1$  divides  $q \pm \sqrt{2q} + 1$ ;
- (c)  $\phi(i)q^2(q^2+1)/2$ , where  $i > 1$  divides  $q-1$ .

*Proof.* Suppose that  $i \in \omega(S)$  is an even number. Then by Lemma 3.1, we have that  $i = 2$  or  $i = 4$  and  $i$  divides the order of subgroup  $W$  as in Lemma 3.2(d). Then each element of  $W$  is of the form  $w(a, b)$  as in (3.1). Obviously,

$$w(a, b)w(c, d) = w(a + c, b + d + (a\pi)c).$$

This in particular shows that  $w(0, b)$  (with  $b \neq 0$ ) are the only elements of  $W$  of order 2. Therefore, the number of involutions in  $W$  is  $q-1$ . Since  $W$  is a part of the partition introduced in Lemma 3.2(e), the elements of order 2 of  $S$  belong to exactly one of the conjugates of  $W$ . Thus by Lemma 3.2(d), there are  $q^2 + 1$  conjugates of  $W$  implying that there are exactly  $m_2(S) = (q-1)(q^2+1)$  involutions in  $S$ . It also follows from Lemma 3.2(d) that the number of elements of order 4 in  $W$  is  $q^2 - q$ , and hence applying the partition in Lemma 3.2(e), we conclude that  $S$  consists of  $m_4(S) = q(q-1)(q^2+1)$  elements of order 4. This proves part (a).

Suppose now  $i \in \omega(S)$  is an odd number. Then, by Lemma 3.1,  $i$  divides the order of one the cyclic subgroups  $U_1, U_2$  and  $V$  as in Lemma 3.2, say  $H$ . Assume  $i = np^\alpha$  with  $p$  odd. Since  $H$  is a part of the partition introduced in Lemma 3.2(e), the elements of order  $i$  are contained in  $H$  and its conjugates. Since also  $H$  is cyclic, there are  $\phi(i)$  elements of order  $i$  in each conjugates of  $H$  including  $H$ .

Now we consider each possibility of  $H$ . If  $H = U_t$  of order  $q \pm \sqrt{2q} + 1$ , for  $t = 1, 2$ , then by Lemma 3.2(b),  $|N_S(U_t) : U_t| = 4$ , and so  $|S : N_S(U_t)| = |S|/4|U_t|$ , for  $t = 1, 2$ . So there are  $|S|/4|U_t|$  conjugates of  $U_t$  implying that there are exactly  $m_i(S) = \phi(i)q^2(q-1)(q \mp \sqrt{2q} + 1)/4$  elements of order  $i$  in  $S$ . This follows part (b). If  $H = V$ , then Lemma 3.2(c) implies that  $|N_S(V) : V| = 2$ , and so the same argument as in the previous cases, we conclude that  $m_i(S) = \phi(i)|S|/2|V| = \phi(i)q^2(q^2+1)/2$ . This follows (c).  $\square$

#### 4. Proof of the Main Theorem

In this section, we prove Theorem 1.1. From now on, set  $S := \text{Sz}(q)$ , where  $q = 2^{2m+1} \geq 8$ , and recall that  $G$  is a finite group with  $\text{nse}(G) = \text{nse}(S)$  and  $|G| = |S|$ . Therefore, by Proposition 3.3,

$nse(S)$  consists of

$$\begin{aligned}
 & m_1(S) = 1; \\
 & m_2(S) = (q - 1)(q^2 + 1); \\
 (4.1) \quad & m_4(S) = q(q - 1)(q^2 + 1); \\
 & m_i(S) = \phi(i)q^2(q \mp \sqrt{2q} + 1)(q - 1)/4, \text{ where } i > 1 \text{ divides } q \pm \sqrt{2q} + 1; \\
 & m_i(S) = \phi(i)q^2(q^2 + 1)/2, \text{ where } i > 1 \text{ divides } q - 1.
 \end{aligned}$$

**Proposition 4.1.** *The vertex 2 is an isolated vertex in  $\Gamma(G)$ .*

*Proof.* Assume the contrary. Then there is an odd prime divisor  $p$  of  $|G|$  such that  $2p \in \omega(G)$ . Let  $f(n)$  be the number of elements of  $G$  whose orders are multiples of  $n$ . Then by Lemma 2.3,  $f(2)$  is a multiple of the greatest divisor of  $|G|$  that is prime to 2. Since  $(q^2 + 1)(q - 1)$  is the greatest divisor of  $|G|$  which is coprime to 2, there exists a positive integer  $r$  such that  $f(2) = (q^2 + 1)(q - 1)r$  and  $(r, 2) = 1$ . On the other hand, by Lemma 2.2, it is obvious that  $m_2(G) = m_2(S)$ , and so

$$f(2) = m_2(G) + \sum_{i>2 \text{ is even}} m_i(G),$$

with  $m_i$  as in (4.1). Now applying Proposition 3.3, there is a non-negative integer  $\alpha$  such that

$$f(2) = (q^2 + 1)(q - 1) + \alpha q(q^2 + 1)(q - 1) + g(2),$$

where

$$g(2) = \sum_{\substack{i|q \pm \sqrt{2q} + 1 \\ i \neq 1}} \beta_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma_i \cdot m_i(S)$$

for some non-negative integers  $\beta_i$  and  $\gamma_i$ . Since  $2p \in \omega(G)$ , we have that  $\alpha q(q^2 + 1)(q - 1) + g(2) > 0$ . Then

$$g(2) = (q^2 + 1)(q - 1)(r - 1 - \alpha q).$$

We now prove that  $q^2$  divides  $g(2)$ . It follows from Lemma 3.1 that 2 is an isolated vertex of  $\Gamma(S)$ . Then a Sylow 2-subgroup of  $S$ , say  $S_2$ , acts fixed point freely (by conjugation) on the set of elements of order  $i \neq 1, 2, 4$  (see Proposition 3.3 and (4.1)). Thus  $|S_2|$  divides  $m_i(S)$  with  $i \neq 1, 2, 4$ . Hence  $q^2$  divides  $m_i(S)$  implying that  $g(2)$  is a multiple of  $q^2$ .

We now consider the following two cases:

(1) Let  $g(2) \neq 0$ . Then  $q^2$  divides  $r - 1 - \alpha q$ , and so  $q^2 + \alpha q + 1 \leq r$ . This implies that  $|G| = q^2(q^2 + 1)(q - 1) < (q^2 + 1)(q - 1)r = f(2)$ , which is impossible.

(2) Let  $g(2) = 0$ . Then  $r - 1 - \alpha q = 0$  and  $\alpha \neq 0$ , and so  $m_{2p}(G) = q(q^2 + 1)(q - 1)$ . Therefore

$$f(p) = \sum_{p|i} m_i(G) = \alpha'q(q^2 + 1)(q - 1) + \sum_{\substack{i|q^2 \pm \sqrt{2}q + 1 \\ i \neq 1}} \beta'_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma'_i \cdot m_i(S),$$

where  $\alpha'$ ,  $\beta'_i$  and  $\gamma'_i$  are non-negative integers. Since  $m_{2p} = q(q^2 + 1)(q - 1)$ , we have that  $\alpha' > 0$ . On the other hand, by Lemma 2.3,  $f(p) = q^2(q^2 + 1)(q - 1)r'/|G_p|$  with  $r'$  a positive integer. Thus

$$(4.2) \quad \frac{q^2(q^2 + 1)(q - 1)r'}{|G_p|} = \alpha'q(q^2 + 1)(q - 1) + \sum_{\substack{i|q^2 \pm \sqrt{2}q + 1 \\ i \neq 1}} \beta'_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma'_i \cdot m_i(S).$$

Since  $q^2$  divides both  $q^2(q^2 + 1)(q - 1)r'/|G_p|$  and  $m_i(S)$  in (4.2), it follows that  $q^2$  divides  $\alpha'q(q^2 + 1)(q - 1)$ . Then  $q \mid \alpha'$ , and so  $|G| = q^2(q^2 + 1)(q - 1) \leq \alpha'q(q^2 + 1)(q - 1) \leq f(p)$ , which is impossible. □

**Proposition 4.2.** *The group  $G$  has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $H$  and  $G/K$  are  $\pi_1$ -groups and  $K/H$  is a non-abelian simple group,  $H$  is a nilpotent group and  $|G/K|$  divides  $|\text{Out}(K/H)|$ .*

*Proof.* By Proposition 4.1, the vertex 2 is an isolated vertex in the prime graph  $\Gamma(G)$  of  $G$ . This implies that the number  $t(G)$  of connected components of the prime graph  $\Gamma(G)$  is at least two. The assertion follows from [16, Theorem A] provided that  $G$  is neither a Frobenius group, nor a 2-Frobenius group.

Let  $G$  be a Frobenius group with kernel  $K$  and complement  $H$ . Then by Lemma 2.4, we must have  $t(G) = 2$ ,  $\pi(H)$  and  $\pi(K)$  are vertex sets of the connected components of  $\Gamma(G)$ . By Proposition 4.1, the vertex 2 is an isolated vertex in  $\Gamma(G)$ . Then either (i)  $|K| = q^2$  and  $|H| = (q^2 + 1)(q - 1)$ , or (ii)  $|H| = q^2$  and  $|K| = (q^2 + 1)(q - 1)$ . Both cases can be ruled out as  $|H|$  must divide  $|K| - 1$ .

Let  $G$  be a 2-Frobenius group. Then Lemma 2.5 implies that  $t(G) = 2$  and  $G$  has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $G/H$  and  $K$  are Frobenius groups with kernel  $K/H$  and  $H$  respectively,  $\pi(G/K) \cup \pi(H) = \pi_1$ ,  $\pi(K/H) = \pi_2$  and  $|G/K|$  divides  $|\text{Aut}(K/H)|$ . Since 2 is an isolated vertex of  $\Gamma(G)$  by Proposition 4.1,  $|K/H| = (q^2 + 1)(q - 1)$  and  $|G/K| \cdot |H| = q^2$ . Since also  $K$  is a Frobenius group with kernel  $H$ , there is a positive integer  $\alpha$  such that  $(q^2 + 1)(q - 1)$  divides  $2^\alpha - 1$ , which is a contradiction. □

**4.1. Proof of Theorem 1.1.**

*Proof.* Let  $S := \text{Sz}(q)$ , where  $q = 2^{2m+1} \geq 8$ . Suppose that  $G$  is a finite group with  $\text{nse}(G) = \text{nse}(S)$  and  $|G| = |S|$ . By applying Proposition 4.2, the group  $G$  has a normal series  $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$  such that  $H$  and  $G/K$  are  $\pi_1$ -groups and  $K/H$  is a non-abelian simple group. Since 3 is a prime divisor of all finite non-abelian simple groups except for Suzuki groups. Moreover, 3 is coprime to  $|K/H|$ . Then  $K/H \cong \text{Sz}(q')$ , where  $q' = 2^{2m'+1}$ . This, in particular, implies that  $2^{4m'+2}$  divides  $2^{4m+2}$ , and hence  $m' \leq m$ . On the other hand,  $H$  and  $G/K$  are  $\pi_1$ -groups. Then  $(q^2 + 1)(q - 1)$  divides  $|K/H|$ ,

and so  $(q^2 + 1)(q - 1)$  divides  $(q'^2 + 1)(q' - 1)$ . Since now  $m' \leq m$ , we must have  $m = m'$ . Therefore  $K/H \cong S$ . Now  $|G| = |K/H| = |S|$ , and hence  $G \cong S$ .  $\square$

### Acknowledgments

The authors are grateful to the anonymous referees and the editor for careful reading of the manuscript and for corrections and suggestions.

### REFERENCES

- [1] S. H. Alavi, A. Daneshkhah and H. P. Mosaed, On quantitative structure of small Ree groups, *Communications in Algebra*, **45** no. 9 (2017) 4099–4108.
- [2] A. Khalili Asboei, Characterization of projective general linear groups, *Int. J. Group Theory*, **5** no. 1 (2016) 17–28.
- [3] A. Khalili Asboei and S. S. Salehi Amiri, A new characterization of  $PSL(2, 25)$ , *Int. J. Group Theory*, **1** no. 3 (2012) 15–19.
- [4] R. W. Carter, *Simple groups of Lie type*, John Wiley & Sons, London-New York-Sydney, **28**, Pure and Applied Mathematics, 1972.
- [5] G. Y. Chen, On structure of Frobenius and 2-Frobenius group, *J. Southwest China Normal Univ.*, **20** no. 5 (1995) 485–487.
- [6] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of finite groups*, Oxford University Press, Eynsham, 1985.
- [7] D. Gorenstein, *Finite groups*, Chelsea Publishing Company., New York, second edition, 1980.
- [8] M. Jr. Hall, *The theory of groups*, The Macmillan Co., New York, N.Y., 1959.
- [9] B. Huppert and N. Blackburn, *Finite groups. III*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **243**, Springer-Verlag, Berlin-New York, 1982.
- [10] A. Iranmanesh, H. Parvizi Mosaed and A. Tehranian, *Characterization of Suzuki group by nse and order of group*, Bulletin of the Korean Mathematical Society, Accepted, 2015.
- [11] V. D. Mazurov and E. I. Khukhro, *Unsolved Problems in Group Theory*, The Kourovka Notebook, **18** 2014, Available at [arxiv.org/abs/1401.0300v6](http://arxiv.org/abs/1401.0300v6).
- [12] L. Mousavi and B. Taeri, A characterization of  $L_2(81)$  by nse, *Int. J. Group Theory*, **5** (1) 29–35 (2016).
- [13] C. Shao, W. Shi and Q. Jiang, Characterization of simple  $K_4$ -groups., *Front. Math. China*, **3** no. 3 (2008) 355–370.
- [14] W. J. Shi, A characterization of Suzuki's simple groups, *Proc. Amer. Math. Soc.*, **114** no. 3 (1992) 589–591.
- [15] L. Weisner, On the number of elements of a group which have a power in a given conjugate set, *Bull. Amer. Math. Soc.*, **31** no. 9-10 (1925) 492–496.
- [16] J. S. Williams, Prime graph components of finite groups, *J. Algebra*, **69** no. 2 (1981) 487–513.

**Seyed Hassan Alavi**

Department of Mathematics, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran

Email: [alavi.s.hassan@basu.ac.ir](mailto:alavi.s.hassan@basu.ac.ir)

Email: [alavi.s.hassan@gmail.com](mailto:alavi.s.hassan@gmail.com)

**Ashraf Daneshkhah**

Department of Mathematics, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran

Email: [adanesh@basu.ac.ir](mailto:adanesh@basu.ac.ir)

Email: [daneshkhah.ashraf@gmail.com](mailto:daneshkhah.ashraf@gmail.com)

**Hosein Parvizi Mosaed**

Alvand Institute of Higher Education, Hamedan, Iran

Email: [h.parvizi.mosaed@gmail.com](mailto:h.parvizi.mosaed@gmail.com)