



www.theoryofgroups.ir



www.ui.ac.ir

DIFFERENCE BASES IN DIHEDRAL GROUPS

TARAS BANAKH AND VOLODYMYR GAVRYLKIV*

Communicated by Attila Maroti

ABSTRACT. A subset B of a group G is called a *difference basis* of G if each element $g \in G$ can be written as the difference $g = ab^{-1}$ of some elements $a, b \in B$. The smallest cardinality $|B|$ of a difference basis $B \subset G$ is called the *difference size* of G and is denoted by $\Delta[G]$. The fraction $\delta[G] := \Delta[G]/\sqrt{|G|}$ is called the *difference characteristic* of G . We prove that for every $n \in \mathbb{N}$ the dihedral group D_{2n} of order $2n$ has the difference characteristic $\sqrt{2} \leq \delta[D_{2n}] \leq \frac{48}{\sqrt{586}} \approx 1.983$. Moreover, if $n \geq 2 \cdot 10^{15}$, then $\delta[D_{2n}] < \frac{4}{\sqrt{6}} \approx 1.633$. Also we calculate the difference sizes and characteristics of all dihedral groups of cardinality ≤ 80 .

1. Introduction

A subset B of a group G is called a *difference basis* for a subset $A \subset G$ if each element $a \in A$ can be written as $a = xy^{-1}$ for some $x, y \in B$. The smallest cardinality of a difference basis for A is called the *difference size* of A and is denoted by $\Delta[A]$. For example, the set $\{0, 1, 4, 6\}$ is a difference basis for the interval $A = [-6, 6] \cap \mathbb{Z}$ witnessing that $\Delta[A] \leq 4$.

The definition of a difference basis B for a set A in a group G implies that $|A| \leq |B|^2$ and gives a lower bound $\sqrt{|A|} \leq \Delta[A]$. The fraction

$$\delta[A] := \frac{\Delta[A]}{\sqrt{|A|}} \geq 1$$

is called the *difference characteristic* of A .

For a real number x we put

$$\lceil x \rceil = \min\{n \in \mathbb{Z} : n \geq x\} \text{ and } \lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}.$$

MSC(2010): Primary: 05B10; Secondary: 05E15, 20D60.

Keywords: dihedral group, difference basis, difference characteristic.

Received: 12 April 2017, Accepted: 08 August 2017.

*Corresponding author.

The following proposition is proved in [3, 1.1].

Proposition 1. *Let G be a finite group. Then*

- (1) $\frac{1+\sqrt{4|G|-3}}{2} \leq \Delta[G] \leq \lceil \frac{|G|+1}{2} \rceil$,
- (2) $\Delta[G] \leq \Delta[H] \cdot \Delta[G/H]$ and $\delta[G] \leq \delta[H] \cdot \delta[G/H]$ for any normal subgroup $H \subset G$;
- (3) $\Delta[G] \leq |H| + |G/H| - 1$ for any subgroup $H \subset G$.

In [10] Kozma and Lev proved (using the classification of finite simple groups) that each finite group G has difference characteristic $\delta[G] \leq \frac{4}{\sqrt{3}} \approx 2.3094$.

In this paper we shall evaluate the difference characteristics of dihedral groups and prove that each dihedral group D_{2n} has $\delta[D_{2n}] \leq \frac{48}{\sqrt{586}} \approx 1.983$. Moreover, if $n \geq 2 \cdot 10^{15}$, then $\delta[D_{2n}] < \frac{4}{\sqrt{6}} \approx 1.633$. We recall that the *dihedral group* D_{2n} is the isometry group of a regular n -gon. The dihedral group D_{2n} contains a normal cyclic subgroup of index 2. A standard model of a cyclic group of order n is the multiplicative group

$$C_n = \{z \in \mathbb{C} : z^n = 1\}$$

of n -th roots of 1. The group C_n is isomorphic to the additive group of the ring $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$.

Difference bases have applications in the study of structure of superextensions of groups, see [1, 3].

A subset B of a group G is called a *basis* of G if each element $g \in G$ can be written as $g = ab$ for some $a, b \in B$. Bases in dihedral groups were studied in [7].

Theorem 2. *For any numbers $n, m \in \mathbb{N}$ the dihedral group D_{2nm} has the difference size*

$$2\sqrt{nm} \leq \Delta[D_{2nm}] \leq \Delta[D_{2n}] \cdot \Delta[C_m]$$

and the difference characteristic $\sqrt{2} \leq \delta[D_{2nm}] \leq \delta[D_{2n}] \cdot \delta[C_m]$.

Proof. It is well-known that the dihedral group D_{2nm} contains a normal cyclic subgroup of order nm , which can be identified with the cyclic group C_{nm} . The subgroup $C_m \subset C_{nm}$ is normal in D_{2nm} and the quotient group D_{2nm}/C_m is isomorphic to D_{2n} . Applying Proposition 1(2), we obtain the upper bounds $\Delta[D_{2n}] \leq \Delta[D_{2nm}/C_m] \cdot \Delta[C_m] = \Delta[D_{2n}] \cdot \Delta[C_m]$ and $\delta[D_{2nm}] \leq \delta[D_{2n}] \cdot \delta[C_m]$.

Next, we prove the lower bound $2\sqrt{nm} \leq \Delta[D_{2nm}]$. Fix any element $s \in D_{2nm} \setminus C_{nm}$ and observe that $s = s^{-1}$ and $sxs^{-1} = x^{-1}$ for all $x \in C_{nm}$. Fix a difference basis $D \subset D_{2nm}$ of cardinality $|D| = \Delta[D_{2nm}]$ and write D as the union $D = A \cup sB$ for some sets $A, B \subset C_{nm} \subset D_{2nm}$. We claim that $AB^{-1} = C_{nm}$. Indeed, for any $x \in C_{nm}$ we get $xs \in sC_{nm} \cap (A \cup sB)(A \cup sB)^{-1} = AB^{-1}s^{-1} \cup sBA^{-1}$ and hence

$$x \in AB^{-1}s^{-1}s^{-1} \cup sBA^{-1}s^{-1} = AB^{-1} \cup B^{-1}A = AB^{-1}.$$

So, $C_{nm} = AB^{-1}$ and hence $nm \leq |A| \cdot |B|$. Then $\Delta[D_{2nm}] = |A| + |B| \geq \min\{l + k : l, k \in \mathbb{N}, lk \geq nm\} \geq 2\sqrt{nm}$ and $\delta[D_{2nm}] = \frac{\Delta[D_{2nm}]}{\sqrt{2nm}} \geq \frac{2\sqrt{nm}}{\sqrt{2nm}} = \sqrt{2}$. □

Corollary 3. *For any number $n \in \mathbb{N}$ the dihedral group D_{2n} has the difference size*

$$2\sqrt{n} \leq \Delta[D_{2n}] \leq 2 \cdot \Delta[C_n]$$

and the difference characteristic $\sqrt{2} \leq \delta[D_{2n}] \leq \sqrt{2} \cdot \delta[C_n]$.

The difference sizes of finite cyclic groups were evaluated in [2] with the help of the difference sizes of the order-intervals $[1, n] \cap \mathbb{Z}$ in the additive group \mathbb{Z} of integer numbers. For a natural number $n \in \mathbb{N}$ by $\Delta[n]$ we shall denote the difference size of the order-interval $[1, n] \cap \mathbb{Z}$ and by $\delta[n] := \frac{\Delta[n]}{\sqrt{n}}$ its difference characteristic. The asymptotics of the sequence $(\delta[n])_{n=1}^\infty$ was studied by Rédei and Rényi [11], Leech [9] and Golay [8] who eventually proved that

$$\sqrt{2 + \frac{4}{3\pi}} < \sqrt{2 + \max_{0 < \varphi < 2\pi} \frac{2 \sin(\varphi)}{\varphi + \pi}} \leq \lim_{n \rightarrow \infty} \delta[n] = \inf_{n \in \mathbb{N}} \delta[n] \leq \delta[6166] = \frac{128}{\sqrt{6166}} < \delta[6] = \sqrt{\frac{8}{3}}.$$

In [2] the difference sizes of the order-intervals $[1, n] \cap \mathbb{Z}$ were applied to give upper bounds for the difference sizes of finite cyclic groups.

Proposition 4. *For every $n \in \mathbb{N}$ the cyclic group C_n has difference size $\Delta[C_n] \leq \Delta[\lceil \frac{n-1}{2} \rceil]$, which implies that*

$$\limsup_{n \rightarrow \infty} \delta[C_n] \leq \frac{1}{\sqrt{2}} \inf_{n \in \mathbb{N}} \delta[n] \leq \frac{64}{\sqrt{3083}} < \frac{2}{\sqrt{3}}.$$

The following upper bound for the difference sizes of cyclic groups were proved in [2].

Theorem 5. *For any $n \in \mathbb{N}$ the cyclic group C_n has the difference characteristic:*

- (1) $\delta[C_n] \leq \delta[C_4] = \frac{3}{2}$;
- (2) $\delta[C_n] \leq \delta[C_2] = \delta[C_8] = \sqrt{2}$ if $n \neq 4$;
- (3) $\delta[C_n] \leq \frac{12}{\sqrt{73}} < \sqrt{2}$ if $n \geq 9$;
- (4) $\delta[C_n] \leq \frac{24}{\sqrt{293}} < \frac{12}{\sqrt{73}}$ if $n \geq 9$ and $n \neq 292$;
- (5) $\delta[C_n] < \frac{2}{\sqrt{3}}$ if $n \geq 2 \cdot 10^{15}$.

For some special numbers n we have more precise upper bounds for $\Delta[C_n]$. A number q is called a *prime power* if $q = p^k$ for some prime number p and some $k \in \mathbb{N}$.

The following theorem was derived in [2] from the classical results of Singer [13], Bose, Chowla [4], [5] and Rusza [12].

Theorem 6. *Let p be a prime number and q be a prime power. Then*

- (1) $\Delta[C_{q^2+q+1}] = q + 1$;
- (2) $\Delta[C_{q^2-1}] \leq q - 1 + \Delta[C_{q-1}] \leq q - 1 + \frac{3}{2}\sqrt{q-1}$;
- (3) $\Delta[C_{p^2-p}] \leq p - 3 + \Delta[C_p] + \Delta[C_{p-1}] \leq p - 3 + \frac{3}{2}(\sqrt{p} + \sqrt{p-1})$.

The following Table 1 of difference sizes and characteristics of cyclic groups C_n for $n \leq 100$ is taken from [2].

TABLE 1. Difference sizes and characteristics of cyclic groups C_n for $n \leq 100$.

n	$\Delta[C_n]$	$\delta[C_n]$	n	$\Delta[C_n]$	$\delta[C_n]$	n	$\Delta[C_n]$	$\delta[C_n]$	n	$\Delta[C_n]$	$\delta[C_n]$
1	1	1	26	6	1.1766...	51	8	1.1202...	76	10	1.1470...
2	2	1.4142...	27	6	1.1547...	52	9	1.2480...	77	10	1.1396...
3	2	1.1547...	28	6	1.1338...	53	9	1.2362...	78	10	1.1322...
4	3	1.5	29	7	1.2998...	54	9	1.2247...	79	10	1.1250...
5	3	1.3416...	30	7	1.2780...	55	9	1.2135...	80	11	1.2298...
6	3	1.2247...	31	6	1.0776...	56	9	1.2026...	81	11	1.2222...
7	3	1.1338...	32	7	1.2374...	57	8	1.0596...	82	11	1.2147...
8	4	1.4142...	33	7	1.2185...	58	9	1.1817...	83	11	1.2074...
9	4	1.3333...	34	7	1.2004...	59	9	1.1717...	84	11	1.2001...
10	4	1.2649...	35	7	1.1832...	60	9	1.1618...	85	11	1.1931...
11	4	1.2060...	36	7	1.1666...	61	9	1.1523...	86	11	1.1861...
12	4	1.1547...	37	7	1.1507...	62	9	1.1430...	87	11	1.1793...
13	4	1.1094...	38	8	1.2977...	63	9	1.1338...	88	11	1.1726...
14	5	1.3363...	39	7	1.1208...	64	9	1.125	89	11	1.1659...
15	5	1.2909...	40	8	1.2649...	65	9	1.1163...	90	11	1.1595...
16	5	1.25	41	8	1.2493...	66	10	1.2309...	91	10	1.0482...
17	5	1.2126...	42	8	1.2344...	67	10	1.2216...	92	11	1.1468...
18	5	1.1785...	43	8	1.2199...	68	10	1.2126...	93	12	1.2443...
19	5	1.1470...	44	8	1.2060...	69	10	1.2038...	94	12	1.2377...
20	6	1.3416...	45	8	1.1925...	70	10	1.1952...	95	12	1.2311...
21	5	1.0910...	46	8	1.1795...	71	10	1.1867...	96	12	1.2247...
22	6	1.2792...	47	8	1.1669...	72	10	1.1785...	97	12	1.2184...
23	6	1.2510...	48	8	1.1547...	73	9	1.0533...	98	12	1.2121...
24	6	1.2247...	49	8	1.1428...	74	10	1.1624...	99	12	1.2060...
25	6	1.2	50	8	1.1313...	75	10	1.1547...	100	12	1.2

Using Theorem 6(1), we shall prove that for infinitely many numbers n the lower and upper bounds given in Theorem 2 uniquely determine the difference size $\Delta[D_{2n}]$ of D_{2n} .

Theorem 7. *If $n = 1 + q + q^2$ for some prime power q , then*

$$\Delta[D_{2n}] = 2 \cdot \Delta[C_n] = \lceil 2\sqrt{n} \rceil = \lceil \sqrt{2|D_{2n}|} \rceil = 2 + 2q.$$

Proof. By Theorem 6(1), $\Delta[C_n] = 1 + q$. Since

$$2\sqrt{q^2 + q + 1} = 2\sqrt{n} \leq \Delta[D_{2n}] \leq \Delta[D_2] \cdot \Delta[C_n] = 2 \cdot \Delta[C_n] = 2 + 2q,$$

it suffices to check that $(2 + 2q) - 2\sqrt{q^2 + q + 1} < 1$, which is equivalent to $\sqrt{q^2 + q + 1} > q + \frac{1}{2}$ and to $q^2 + q + 1 > q^2 + q + \frac{1}{4}$. \square

A bit weaker result holds also for the dihedral groups $D_{8(q^2+q+1)}$.

Proposition 8. *If $n = 1 + q + q^2$ for some prime power q , then*

$$4q + 3 \leq \Delta[D_{8n}] \leq 4q + 4.$$

Proof. By Theorem 6(1), $\Delta[C_n] = 1 + q$. Since $\Delta[D_8] = 4$ (see Table 2), by Theorem 2,

$$4\sqrt{q^2 + q + 1} = 2\sqrt{4n} \leq \Delta[D_{8n}] \leq \Delta[D_8] \cdot \Delta[C_n] = 4(1 + q).$$

To see that $4q + 3 \leq \Delta[D_{8n}] \leq 4q + 4$, it suffices to check that $(4 + 4q) - 4\sqrt{q^2 + q + 1} < 2$, which is equivalent to $\sqrt{q^2 + q + 1} > q + \frac{1}{2}$ and to $q^2 + q + 1 > q^2 + q + \frac{1}{4}$. \square

In Table 2 we present the results of computer calculation of the difference sizes and characteristics of dihedral groups of order ≤ 80 . In this table $lb[D_{2n}] := \lceil \sqrt{4n} \rceil$ is the lower bound given in Theorem 2. With the boldface font we denote the numbers $2n \in \{14, 26, 42, 62\}$, equal to $2(q^2 + q + 1)$ for a prime power q . For these numbers we know that $\Delta[D_{2n}] = lb[D_{2n}] = 2q + 2$. For $q = 2$ and $n = q^2 + q + 1 = 7$ the table shows that $\Delta[D_{56}] = \Delta[D_{8n}] = 11 = 4q + 3$, which means that the lower bound $4q + 3$ in Proposition 8 is attained.

Theorem 9. *For any number $n \in \mathbb{N}$ the dihedral group D_{2n} has the difference characteristic*

$$\sqrt{2} \leq \delta[D_{2n}] \leq \frac{48}{\sqrt{586}} \approx 1.983.$$

Moreover, if $n \geq 2 \cdot 10^{15}$, then $\delta[D_{2n}] < \frac{4}{\sqrt{6}} \approx 1.633$.

Proof. By Corollary 3, $\sqrt{2} \leq \delta[D_{2n}] \leq \sqrt{2} \cdot \delta[C_n]$. If $n \geq 9$ and $n \neq 292$, then $\delta[C_n] \leq \frac{24}{\sqrt{293}}$ by Theorem 5(4), and hence $\delta[D_{2n}] \leq \sqrt{2} \cdot \delta[C_n] \leq \sqrt{2} \cdot \frac{24}{\sqrt{293}} = \frac{48}{\sqrt{586}}$. If $n = 292$, then known values $\delta[C_{73}] = \frac{9}{\sqrt{73}}$ (given in Table 1), $\delta[D_8] = \frac{4}{\sqrt{8}} = \sqrt{2}$ (given in Table 2) and Theorem 2 yield the upper bound

$$\delta[D_{2 \cdot 292}] = \delta[D_{8 \cdot 73}] \leq \delta[D_8] \cdot \delta[C_{73}] = \sqrt{2} \cdot \frac{9}{\sqrt{73}} < \frac{48}{\sqrt{586}}.$$

Analyzing the data from Table 2, one can check that $\delta[D_{2n}] \leq \frac{48}{\sqrt{586}} \approx 1.983$ for all $n \leq 8$.

If $n \geq 2 \cdot 10^{15}$, then $\delta[C_n] < \frac{2}{\sqrt{3}}$ by Theorem 5(5), and hence

$$\delta[D_{2n}] \leq \sqrt{2} \cdot \delta[C_n] < \frac{4}{\sqrt{6}}.$$

\square

Question 10. *Is $\sup_{n \in \mathbb{N}} \delta[D_{2n}] = \delta[D_{22}] = \frac{8}{\sqrt{22}} \approx 1.7056$?*

To answer Question 10 affirmatively, it suffices to check that $\delta[D_{2n}] \leq \frac{8}{\sqrt{22}}$ for all $n < 1\,212\,464$.

TABLE 2. Difference sizes and characteristics of dihedral groups D_{2n} for $2n \leq 80$.

$2n$	$lb[D_{2n}]$	$\Delta[D_{2n}]$	$2\Delta[C_n]$	$\delta[D_{2n}]$	$2n$	$lb[D_{2n}]$	$\Delta[D_{2n}]$	$2\Delta[C_n]$	$\delta[D_{2n}]$
2	2	2	2	1.4142...	42	10	10	10	1.5430...
4	3	3	4	1.5	44	10	10	12	1.5075...
6	4	4	4	1.6329...	46	10	11	12	1.6218...
8	4	4	6	1.4142...	48	10	10	12	1.4433...
10	5	5	6	1.5811...	50	10	11	12	1.5556...
12	5	5	6	1.4433...	52	11	11	12	1.5254...
14	6	6	6	1.6035...	54	11	12	12	1.6329...
16	6	6	8	1.5	56	11	11	12	1.4699...
18	6	7	8	1.6499...	58	11	12	14	1.5756...
20	7	7	8	1.5652...	60	11	12	14	1.5491...
22	7	8	8	1.7056...	62	12	12	12	1.5240...
24	7	7	8	1.4288...	64	12	12	14	1.5
26	8	8	8	1.5689...	66	12	13	14	1.6001...
28	8	8	10	1.5118...	68	12	13	14	1.5764...
30	8	8	10	1.4605...	70	12	12	14	1.4342...
32	8	9	10	1.5909...	72	12	13	14	1.5320...
34	9	9	10	1.5434...	74	13	14	14	1.6274...
36	9	9	10	1.5	76	13	14	16	1.6059...
38	9	10	10	1.6222...	78	13	14	14	1.5851...
40	9	9	12	1.4230...	80	13	14	16	1.5652...

Proposition 11. The inequality $\delta[D_{2n}] \leq \sqrt{2} \cdot \delta[C_n] \leq \frac{8}{\sqrt{22}}$ holds for all $n \geq 1\,212\,464$.

Proof. It suffices to prove that $\delta[C_n] \leq \frac{4}{\sqrt{11}}$ for all $n \geq 1\,212\,464$. To derive a contradiction, assume that $\delta[C_n] > \frac{4}{\sqrt{11}}$ for some $n \geq 1\,212\,464$. Let $(q_k)_{k=1}^\infty$ be an increasing enumeration of prime powers. Let $k \in \mathbb{N}$ be the unique number such that $12q_k^2 + 14q_k + 15 < n \leq 12q_{k+1}^2 + 14q_{k+1} + 15$. By Corollary 4.9 of [2], $\Delta[C_n] \leq 4(q_{k+1} + 1)$. The inequality $\delta[C_n] > \frac{4}{\sqrt{11}}$ implies

$$4(q_{k+1} + 1) \geq \Delta[C_n] > \frac{4}{\sqrt{11}}\sqrt{n} \geq \frac{4}{\sqrt{11}}\sqrt{12q_k^2 + 14q_k + 16}.$$

By Theorem 1.9 of [6], if $q_k \geq 3275$, then $q_{k+1} \leq q_k + \frac{q_k}{2\ln^2(q_k)}$. On the other hand, using WolframAlpha computational knowledge engine it can be shown that the inequality $1 + x + \frac{x}{2\ln^2(x)} \leq \frac{1}{\sqrt{11}}\sqrt{12x^2 + 14x + 16}$ holds for all $x \geq 43$. This implies that $q_k < 3275$.

Analyzing the table¹ of (maximal gaps between) primes, it can be shown that $11(q_{k+1} + 1)^2 \leq 12q_k^2 + 14q_k + 16$ if $q_k \geq 331$. So, $q_k \leq 317$, $q_{k+1} \leq 331$ and $11 \cdot (q_{k+1} + 1)^2 = 11 \cdot 332^2 = 1\,212\,464 \leq n$, which contradicts $4(q_{k+1} + 1) > \frac{4}{\sqrt{11}}\sqrt{n}$. \square

REFERENCES

- [1] T. Banakh and V. Gavrylkiv, Algebra in the superextensions of twinic groups, *Dissertationes Math.* **473** (2010) pp. 74.
- [2] T. Banakh and V. Gavrylkiv, Difference bases in cyclic groups, *em J. Algebra Appl.* **18** (2019) pp. 18.
- [3] T. Banakh, V. Gavrylkiv and O. Nykyforchyn, Algebra in superextension of groups, I: zeros and commutativity, *Algebra Discrete Math.*, **3** (2008) 1–29.
- [4] R. C. Bose, An affine analogue of Singer's theorem, *J. Indian Math. Soc.*, **6** (1942) 1–15.
- [5] R. C. Bose and S. Chowla, Theorems in the additive theory of numbers, *Comment. Math. Helv.*, **37** (1962-63) 141–147.
- [6] P. Dusart, *Autour de la fonction qui compte le nombre de nombres premiers*, Ph.D. Thesis, Univ. de Limoges, 1998.
- [7] V. Gavrylkiv, Bases in dihedral and Boolean groups, *J. Integer Seq.*, **20** (2017).
- [8] M. Golay, Notes on the representation of $1, 2, \dots, n$ by differences, *J. Lond. Math. Soc.*, **4** (1972) 729–734.
- [9] J. Leech, On the representation of $1, 2, \dots, n$ by differences, *J. Lond. Math. Soc.*, **31** (1956) 160–169.
- [10] G. Kozma and A. Lev, Bases and decomposition numbers of finite groups, *Arch. Math. (Basel)*, **58** (1992) 417–424.
- [11] L. Rédei and A. Rényi, On the representation of the numbers $1, 2, \dots, N$ by means of differences, *Math. Sbornik N.S.*, **24** (1949) 385–389.
- [12] I. Z. Ruzsa, Solving a linear equation in a set of integers I, *Acta Arith.*, **3** (1993) 259–282.
- [13] J. Singer, A theorem in finite projective geometry and some applications to number theory, *Trans. Amer. Math. Soc.*, **43** (1938) 377–385.

Taras Banakh

Ivan Franko National University of Lviv (Ukraine), and
 Institute of Mathematics, Jan Kochanowski University in Kielce (Poland)
 Email: t.o.banakh@gmail.com

Volodymyr Gavrylkiv

Vasyl Stefanyk Precarpathian National University, Ivano-Frankivsk, Ukraine
 Email: vgavrylkiv@gmail.com

¹See <https://primes.utm.edu/notes/GapsTable.html> and <https://primes.utm.edu/lists/small/1000.txt>