



www.theoryofgroups.ir

International Journal of Group Theory
ISSN (print): 2251-7650, ISSN (on-line): 2251-7669
Vol. 8 No. 2 (2019), pp. 25-28.
© 2019 University of Isfahan



www.ui.ac.ir

B_π -CHARACTERS AND QUOTIENTS

MARK L. LEWIS

Communicated by Attila Maroti

ABSTRACT. Let π be a set of primes, and let G be a finite π -separable group. We consider the Isaacs B_π -characters. We show that if N is a normal subgroup of G , then $B_\pi(G/N) = \text{Irr}(G/N) \cap B_\pi(G)$.

1. Introduction

All groups in this paper are finite. Throughout this paper π will be a set of primes and G will be a π -separable group. In [3], Isaacs defined the subset $B_\pi(G)$ of $\text{Irr}(G)$. In our paper [6] about a variation on Landau's theorem, we needed a basic fact about the characters in $B_\pi(G)$ and quotients of G that seems not to have been proved anywhere. Since proving this fact in [6] would have been a distraction to the main point of that paper, we decided to establish the proof of this fact separately.

Many of the basic ideas about the set $B_\pi(G)$ were proved in [3], and in fact, all of the facts we need about $B_\pi(G)$ can be found in [3]. The papers [4] and [5] both give very good expository accounts about $B_\pi(G)$ characters. Much of this paper is to provide the terminology, concepts, and definitions needed to explicitly define the set $B_\pi(G)$.

Suppose G is a group and N is a normal subgroup of G . There is a bijection between the sets $\{\chi \in \text{Irr}(G) \mid N \leq \ker(\chi)\}$ and $\text{Irr}(G/N)$ (see Lemma 2.22 of [2]). If $\chi \in \text{Irr}(G)$ and $N \leq \ker(\chi)$, then we write $\hat{\chi}$ to denote the corresponding character in $\text{Irr}(G/N)$. With this in mind, we set the

MSC(2010): Primary: 20C15.

Keywords: B_π -characters, π -theory, π -separable groups.

Received: 10 August 2017, Accepted: 22 November 2017.

<http://dx.doi.org/10.22108/ijgt.2017.105879.1442>

following notation: $\hat{B}_\pi(G) = \{\hat{\chi} \mid \chi \in B_\pi(G), N \leq \ker(\chi)\}$. Notice that if we identify $\text{Irr}(G/N)$ and $B_\pi(G/N)$ with the appropriate subsets of $\text{Irr}(G)$, then $\hat{B}_\pi(G) = \text{Irr}(G/N) \cap B_\pi(G)$.

The goal of this paper is to prove the following:

Theorem 1.1. *Suppose π is a set of primes and G is a π -separable group. If N is a normal subgroup of G , then $B_\pi(G/N) = \hat{B}_\pi(G)$.*

2. Results

The first idea that we need for defining the set $B_\pi(G)$ are π -special characters. Following Gajendragadkar in [1], we say that a character $\chi \in \text{Irr}(G)$ where G is π -special if $\chi(1)$ is a π -number and for every subnormal subgroup S of G , the irreducible constituents of χ_S have determinantal order that is a π -number.

In Proposition 7.1 of [1], Gajendragadkar proved that if $\alpha, \beta \in \text{Irr}(G)$ are characters so that α is π -special and β is π' -special, then $\alpha\beta$ is irreducible, and this factorization is unique. I.e., if $\alpha\beta = \alpha'\beta'$ where α' is π -special and β' is π' -special, then $\alpha = \alpha'$ and $\beta = \beta'$.

Using [3], we say that $\chi \in \text{Irr}(G)$ is π -factored if there exists a π -special character α and π' -special character β so that $\chi = \alpha\beta$. We take time out of our definitions to prove the following lemma regarding the kernels of π -factored characters which is key to our argument.

Lemma 2.1. *Suppose π is a set of primes and G is a π -separable group. If $\chi \in \text{Irr}(G)$ satisfies $\chi = \alpha\beta$ where α is π -special and β is π' -special, then $\ker(\chi) = \ker(\alpha) \cap \ker(\beta)$.*

Proof. It is obvious that $\ker(\alpha) \cap \ker(\beta) \leq \ker(\chi) = K$. We need to show that $K \leq \ker(\alpha) \cap \ker(\beta)$. We first claim that $K \leq Z(\alpha) \cap Z(\beta)$. Suppose $g \in K$. Then $\alpha(g)\beta(g) = \chi(g) = \chi(1) = \alpha(1)\beta(1)$. Hence, $\alpha(1)\beta(1) = |\alpha(g)\beta(g)| = |\alpha(g)||\beta(g)|$. By Lemma 2.15(c) of [2], we know that $|\alpha(g)| \leq \alpha(1)$ and $|\beta(g)| \leq \beta(1)$. The previous equality implies that these inequalities must be equalities, so $g \in Z(\alpha)$ and $g \in Z(\beta)$. This proves the claim.

By Lemma 2.27 (c) of [2], we see that $\alpha_K = \alpha(1)\mu$ and $\beta_K = \beta(1)\nu$ for linear characters μ and ν in $\text{Irr}(K)$. Because α is π -special, μ must have π -order and because β is π' -special, ν must have π' -order. For $g \in K$, this implies that $\mu(g)$ is a π root of unity and $\nu(g)$ is a π' -root of unity. We have $\alpha(1)\beta(1) = \chi(1) = \chi(g) = \alpha(g)\beta(g) = \alpha(1)\mu(g)\beta(1)\nu(g)$. This implies that $\nu(g)\mu(g) = 1$. The only way that the product of a π -root of unity and a π' -root can equal 1 is if they are both 1. I.e., we must have $\mu(g) = \nu(g) = 1$. This implies that $\alpha(1) = \alpha(g)$ and $\beta(1) = \beta(g)$. Therefore, $g \in \ker(\alpha) \cap \ker(\beta)$ as desired. \square

Returning to our definitions, we fix the character $\chi \in \text{Irr}(G)$. We say that (S, σ) is a *subnormal pair* for χ if S is a subnormal subgroup of G and σ is an irreducible constituent of χ_S . In addition, we say that (S, σ) is π -factored if σ is π -factored. We can define a partial ordering on the subnormal pairs for χ by $(S, \sigma) \leq (T, \tau)$ if $S \leq T$ and σ is a constituent of τ_S .

Notice that $(1, 1_1)$ is a π -factored subnormal pair for χ , so there exists a maximal π -factored subnormal pair for χ with respect to the partial ordering. It is shown in Theorem 3.2 of [3] that

the set of maximal π -factored subnormal pairs for χ are conjugate in G . Let (S, σ) be a maximal π -factored subnormal pair for χ , and let T be the stabilizer of (S, σ) in G . It is shown in Theorem 4.4 of [3] that there is a unique character $\tau \in \text{Irr}(T \mid \sigma)$ so that $\tau^G = \chi$.

We can now define the π -nucleus for χ . If χ is π -factored, then (G, χ) is the nucleus for χ . If χ is not π -factored, then let (S, σ) be a maximal π -factored subnormal pair for χ . Let T be the stabilizer of (S, σ) in G , and let $\tau \in \text{Irr}(T \mid \sigma)$ so that $\tau^G = \chi$. By Lemma 4.5 of [3], we know that $T < G$, so we can inductively define the π -nucleus of χ to be the π -nucleus of τ . Because the maximal π -factored subnormal pairs are all conjugate, it follows that the π -nucleus for χ is well-defined up to conjugacy. (See the argument on page 108 of [3].)

We are now ready to state the definition of $B_\pi(G)$. We still have the character χ . We take (X, η) to be a π -nucleus for χ , and by definition η must be π -factored. The set $B_\pi(G)$ is defined to be those characters $\chi \in \text{Irr}(G)$ where (X, η) is a π -nucleus for χ and η is π -special. This is the statement of Definition 5.1 of [3].

We now give a lemma that connects a π -nucleus with the quotient.

Lemma 2.2. *Let π be a set of primes and let G be a π -separable group. Suppose that N is a normal subgroup of G . If $\chi \in \text{Irr}(G)$ with $N \leq \ker(\chi)$ has π -nucleus (X, η) , then $(X/N, \hat{\eta})$ is a π -nucleus for $\hat{\chi} \in \text{Irr}(G/N)$.*

Proof. If $(X, \eta) = (G, \chi)$, then this is obvious. Thus, we may assume that $X < G$. Let (S, σ) be a maximal π -factored subnormal pair for χ with stabilizer T and character $\tau \in \text{Irr}(T \mid \sigma)$ so that $\tau^G = \chi$ and (X, η) is a π -nucleus for τ . Notice that $(N, 1_N)$ is a π -factored subnormal pair for χ , so it is contained in a maximal such pair. As we mentioned above, the maximal π -factored subnormal pairs for χ are all conjugate. Since N is normal, this implies that $N \leq S$, and thus, $(N, 1_N) \leq (S, \sigma)$. Because σ is a constituent of χ_S , we see that $N \leq \ker(\sigma)$. By Lemma 2.1, we see that $\hat{\sigma}$ is π -factored as a character in $\text{Irr}(S/N)$. Notice that $(S/N, \hat{\sigma}) \leq (S^*/N, \hat{\sigma}^*)$ if and only if $(S, \sigma) \leq (S^*, \sigma^*)$, and by Lemma 2.1, $\hat{\sigma}^*$ is π -factored in $\text{Irr}(S^*/N)$ if and only if σ is π -factored in $\text{Irr}(S^*)$. Therefore, $(S/N, \hat{\sigma})$ must be a maximal π -factored subnormal pair for $\hat{\chi}$. It is immediate that T/N will be the stabilizer for $(S/N, \hat{\sigma})$ in G/N and that $\hat{\tau}$ is the unique character in $\text{Irr}(T/N \mid \hat{\sigma})$ that induces $\hat{\chi}$. By induction, $(X/N, \hat{\eta})$ will be the π -nucleus for $\hat{\tau}$, and thus, $(X/N, \hat{\eta})$ will be a π -nucleus for $\hat{\chi}$. □

We are now ready to prove Theorem 1.1.

Proof of Theorem 1.1. Note that $B_\pi(G/N) \subseteq \text{Irr}(G/N)$. Hence, it suffices to show for $\chi \in \text{Irr}(G)$ with $N \leq \ker(\chi)$ that $\chi \in B_\pi(G)$ if and only if $\hat{\chi} \in B_\pi(G/N)$. Suppose $\chi \in \text{Irr}(G/N)$. Let (X, η) be a π -nucleus for χ . By Lemma 2.2, $(X/N, \hat{\eta})$ is a nucleus for $\hat{\chi}$. We know that $\chi \in B_\pi(G)$ if and only if η is π -special and $\hat{\chi} \in B_\pi(G/N)$ if and only if $\hat{\eta}$ is π -special. Using the definition of π -special, we see that η is π -special if and only if $\hat{\eta}$ is π -special, and this proves the theorem. □

REFERENCES

- [1] D. Gajendragadkar, A characteristic class of characters of finite p -separable groups, *J. Algebra*, **59** (1979) 237–259.
- [2] I. M. Isaacs, *Character Theory of Finite Groups*, Academic Press, San Diego, California, 1976.
- [3] I. M. Isaacs, Characters of π -separable groups, *J. Algebra* **86** (1984) 98–128.
- [4] I. M. Isaacs, The π -character theory of solvable groups, *J. Austral. Math. Soc. (Series A)*, **57** (1994) 81–102.
- [5] I. M. Isaacs, Characters and sets of primes for solvable groups, in *Finite and locally finite groups (Istanbul, 1994)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., **471**, Kluwer Acad. Publ., Dordrecht, 1995 347–376.
- [6] M. L. Lewis, Landau’s theorem, fields of values for characters, and solvable groups, *J. Austral. Math. Soc.*, (2016) 1–7.

Mark L. Lewis

Department of Mathematical Sciences, Kent State University, Kent, Ohio, USA

Email: lewis@math.kent.edu