



www.theoryofgroups.ir

International Journal of Group Theory
ISSN (print): 2251-7650, ISSN (on-line): 2251-7669
Vol. 2 No. 1 (2013), pp. 157-166.
© 2013 University of Isfahan



www.ui.ac.ir

SUPERSOLUBLE CONDITIONS AND TRANSFER CONTROL

A. L. GILOTTI AND L. SERENA*

Communicated by Patrizia Longobardi

ABSTRACT. In this paper we give a new condition for a Sylow p -subgroup of a finite group to control transfer. Then it is deduced a characterization of supersoluble groups that can be seen as a generalization of the well known result concerning the supersolubility of finite groups with cyclic Sylow subgroups. Moreover a condition for a normal embedding of a strongly closed p -subgroup is given. These results make use of the properties of G -chains and Φ -chains.

1. Introduction

Let G be a finite group, P be a Sylow p -subgroup of G and $V \trianglelefteq P$. Then we say that V controls (strong) fusion in P with respect to G if for any $g \in G$ and any subset A of P such that $A^g \subseteq P$ then $\exists t \in N_G(V)$ such that $A^g = A^t$ (if it controls strong fusion then $g = ct$ with $c \in C_G(A)$ and $t \in N_G(V)$). Moreover we say that V controls transfer in P with respect to G if $P \cap G' = P \cap N_G(V)'$. It follows by the definitions and by the Focal Subgroup Theorem (see [16, p. 142]) that if V controls fusion, then it controls transfer.

These concepts play a fundamental role in the study of local properties of a finite group. We observe that some authors prefer to say that $N_G(V)$, instead of V , controls (strong) fusion or controls transfer.

In the papers [9] and [10] two types of chains have been introduced, namely the G -chain and the Φ -chain of a strongly closed subgroup whose definitions will be recalled below. These chains play a role, respectively, for the control of strong fusion and of transfer by a strongly closed subgroup. In particular, conditions for the existence of a normal Sylow p -complement can be deduced from their properties. In this paper we improve some criterions of [10] for the control of transfer from which we deduce

MSC(2010): Primary: 20D20; Secondary: 20D25, 20D99, 20F16.

Keywords: normal Sylow p -complement, supersoluble groups, strongly closed subgroup, transfer control.

Received: 27 June 2012, Accepted: 14 March 2013.

*Corresponding author.

new conditions for the existence of normal Sylow p -complements and the following characterization of supersoluble groups which can be seen as a generalization of the well known result concerning the supersolubility of finite groups with cyclic Sylow subgroups:

A finite group G is supersoluble if and only if there is a normal subgroup N such that G/N has cyclic Sylow p -subgroups and for every Sylow subgroup P of G we have that $P \cap N$ has a Φ -chain.

Using properties of G -chains and following some ideas from [10] we prove the supersolubility of a finite group G in which all cyclic subgroups of order p (for all primes p which divide $|G|$) or 4 are strongly closed. P. Csörgö and M. Herzog proved the same result, with different proof and by using the concept of \mathcal{H} -subgroup (see [3, Th. 8]). We observe that for a p -subgroup, the two concepts of strongly closed subgroup and of \mathcal{H} -subgroup are equivalent (see Lemma 2.2).

If G is p -soluble, then a p -subgroup V which is strongly closed can be characterized as the Sylow p -subgroup of its normal closure. A subgroup possessing this property is called normally embedded (see for instance [4, p. 250]). The above condition is not longer true in general. However, using properties of G -chains we can prove the following result:

Let G be a finite group, $p \in \pi(G)$ and $P \in \text{Syl}_p(G)$. Let V be strongly closed in P and suppose that there is a chain $1 = V_0 \triangleleft V_1 \dots \triangleleft V_n = V$ with V_i , $i = 1, \dots, n$ weakly closed and $|V_i : V_{i-1}| = p$. Moreover, suppose that $(|N_G(V) : C_G(V)|, p - 1) = 1$, then $V \in \text{Syl}_p(V^G)$.

2. Some definitions and preliminary results

All the groups considered in the paper are finite and the notation is usually standard. In particular $\pi(G)$ indicates the set of different primes which divide the order of a finite group G . If $p \in \pi(G)$ we set $p' = \{\pi(G) \setminus p\}$. If G is the semidirect product of the subgroups H and K with $H \trianglelefteq G$, we write $[H]K$. If $p \in \pi(G)$, then $\text{Syl}_p(G)$ is the set of all the Sylow p -subgroups of G . A finite group G is said to possess a Sylow tower if there is a normal series $1 = G_0 \triangleleft G_1 \triangleleft G_2 \dots \triangleleft G_n = G$ in G such that G_i/G_{i-1} , $1 \leq i \leq n$ is isomorphic to a Sylow subgroup of G . $O_p(G)$ is the maximal normal p -subgroup of G and $O_{p'}(G)$ is the maximal normal subgroup whose order is not divisible by p . $O^p(G)$ indicates the minimal normal subgroup of G such that $G/O^p(G)$ is a p -group. If $O^p(G) = O_{p'}(G)$, then this is a normal Sylow p -complement of G and G is said to be p -nilpotent. If A is a subgroup of a finite group G , then $N_G(A)$ and $C_G(A)$ are respectively the normalizer and the centralizer of A in G . Moreover A^G is the normal closure of A i.e. the minimal normal subgroup of G containing A . Finally, if G is a p -group, then $\Omega_i(G)$ is the subgroup of G generated by the elements of G of order less or equal to p^i .

For the sake of the reader we remember some results and definitions.

Definition 2.1. *Let T be a subgroup of a finite group G and let A be a subset of T .*

- a) *A is said to be weakly closed in T (with respect to G) if whenever $A^g \subseteq T$ with $g \in G$, we have $A^g = A$.*
- b) *A is said to be strongly closed in T (with respect to G) if for any element $a \in A$ and for any $g \in G$, $a^g \in T$ implies $a^g \in A$ (that is $A^g \cap T \subseteq A$).*

- c) If A is a subgroup and $T = N_G(A)$, then A is said to be an \mathcal{H} -subgroup of G if $A^g \cap T \leq A$ for all $g \in G$ (see [3]).

We often will omit the "with respect to G " if it is not necessary to specify in which group we consider the property.

We have the following properties of strongly closed subgroups (see [16, pg. 584] and [10]).

Lemma 2.2. *Let G be a finite group. Then*

- (1) *If A is a strongly closed subset in a subgroup T of G with respect to G , then A is weakly closed in T with respect to G .*
- (2) *Let V be a p -subgroup of G and let $P \in \text{Syl}_p(G)$ such that $V \leq P$. Then the following conditions on V are equivalent:*
 - *The subgroup V is an \mathcal{H} -subgroup of G .*
 - *If S is any p -subgroup of G such that $V \leq S$, then V is strongly closed in S with respect to G .*
 - *The subgroup V is strongly closed in P with respect to G .*
- (3) *Let V be a p -subgroup of G and let $P \in \text{Syl}_p(G)$ such that $V \leq P$. Suppose that V is a strongly closed subgroup in P with respect to G . Then the following properties hold:*
 - i) *For any normal subgroup N of G , $V \cap N$ is a strongly closed subgroup in P with respect to G .*
 - ii) *For any normal subgroup N of G , let $\bar{G} = G/N$. Then $\bar{V} = VN/N$ is a strongly closed subgroup in $\bar{P} = PN/N$ with respect to \bar{G} and we have*

$$N_{\bar{G}}(\bar{V}) = N_G(V)N/N.$$

- (4) *Let G be a finite group and $P \in \text{Syl}_p(G)$. Suppose that $V \trianglelefteq P$. Then*
 - a) *If $N \trianglelefteq G$ and V is weakly closed in P with respect to G then VN/N is weakly closed in PN/N with respect to G/N .*
 - b) *V is normal in G if and only if V is weakly closed in P and subnormal in G .*

Now we summarize and generalize some definitions given in previous papers (see [8] and [10]).

Definitions 2.3. a) *Let G be a finite group and $P \in \text{Syl}_p(G)$. Suppose that V_a and V_b are strongly closed subgroups of P with $V_a < V_b$. Let*

$$(2.1) \quad V_a = V_o \trianglelefteq V_1 \trianglelefteq \dots \trianglelefteq V_n = V_b$$

a chain where $V_i, i = 1, \dots, n - 1$ is weakly closed in P with respect to G and $V_i/V_{i-1} \leq Z(V_b/V_{i-1})$ for $i = 1, \dots, n$. Then (2.1) is said to be a $p - G$ -chain connecting V_a and V_b (or, if there is no ambiguity, a G -chain connecting V_a and V_b). If $V_a = 1$ we simply say that (2.1) is a G -chain of V_b .

- b) *If in the above definition $|V_i : V_{i-1}| = p$ ($i = 1, \dots, n$), then (2.1) is said to be a strict $p - G$ -chain connecting V_a and V_b .*

- c) Suppose that V is a strongly closed subgroup of $P \in \text{Syl}_p(G)$ and that $\Phi(V)$ is also strongly closed in P . Then a strict G -chain connecting $\Phi(V)$ and V is said to be a Φ -chain of V , that is

$$\Phi(V) = V_0 \trianglelefteq V_1 \trianglelefteq \dots \trianglelefteq V_n = V$$

with V_i weakly closed and $|V_i/V_{i-1}| = p$ ($i = 1, \dots, n$).

3. Supersoluble conditions

Let V be a normal subgroup of a Sylow p -subgroup P of a finite group G . Then V controls transfer, i.e. $P \cap G' = P \cap N_G(V)'$ if and only if $G/O^p(G)G' \cong N_G(V)/(O^p(N_G(V))N_G(V)')$ ([14, Lemma 6.15 d), p. 49]). On the other hand $G/O^p(G)G' \cong N_G(V)/(O^p(N_G(V))N_G(V)')$ if and only if $G/O^p(G) \cong N_G(V)/O^p(N_G(V))$ (see for instance [14, Th. 6.18, p. 51]). So, in the following, when we say that V controls transfer we mean

$$G/O^p(G) \cong N_G(V)/O^p(N_G(V)).$$

Lemma 3.1. [10], Lemma 21] Let G be a finite group and $P \in \text{Syl}_p(G)$. Suppose that V is a strongly closed subgroup of P . Moreover suppose that $\Phi(V)$ is strongly closed in P with respect to G . Then V controls transfer.

Proposition 3.2. Let G be a finite group and P a Sylow p -subgroup of G . Suppose that V is a strongly closed subgroup of P which possesses a Φ -chain and that P/V is cyclic. Then P controls transfer.

Proof. Suppose that G is a minimal counterexample and let $N = N_G(P)$ and $N_1 = N_G(V)$. Since V possesses a Φ -chain, by Lemma 3.1 we have $G/O^p(G) \cong N_1/O^p(N_1)$. If $N_1 < G$, the hypotheses go to N_1 and so P controls transfer in N_1 , that is

$$N_{N_1}(P)/O^p(N_{N_1}(P)) \cong N_1/O^p(N_1).$$

Since $N_G(P)/O^p(N_G(P)) = N_{N_1}(P)/O^p(N_{N_1}(P))$ we have

$$G/O^p(G) \cong N_G(P)/O^p(N_G(P)).$$

Therefore we can assume that $N_1 = G$, that is $V \trianglelefteq G$. Then we have $\Phi(V) \trianglelefteq G$ and $\Phi(V) \leq \Phi(P)$. Suppose that $\Phi(V) \neq 1$ and consider $\widehat{G} = G/\Phi(V)$. Then $N/\Phi(V) = N_{\widehat{G}}(P/\Phi(V))$. By minimality of G we have $G/\Phi(V)/O^p(G/\Phi(V)) \cong N/\Phi(V)/O^p(N/\Phi(V))$ that is

$$G/O^p(G)\Phi(V) \cong N/O^p(N)\Phi(V).$$

Then $P \cap O^p(G)\Phi(V) = P \cap O^p(N)\Phi(V)$ that is $\Phi(V)(P \cap O^p(G)) = \Phi(V)(P \cap O^p(N))$. Since $\Phi(V) \leq \Phi(P)$ it follows that $\Phi(P)(P \cap O^p(G)) = \Phi(P)(P \cap O^p(N))$. We have that $G = PO^p(G)$ and $N = PO^p(N)$. So it follows that $\Phi(P)O^p(G) \trianglelefteq G$ and $\Phi(P)O^p(N) \trianglelefteq N$. On the other hand $P/(P \cap O^p(G)\Phi(P))$ is elementary abelian as well $P/(P \cap O^p(N)\Phi(P))$. Therefore $O^p(G)\Phi(P) \geq G'$ and then $O^p(G)\Phi(P) = O^p(G)G'\Phi(P)$. Similarly $O^p(N)\Phi(P) \geq N'$ and so $O^p(N)\Phi(P) = O^p(N)N'\Phi(P)$. In particular $O^p(G)\Phi(P)$ and $O^p(N)\Phi(P)$ are the smallest normal subgroups of G and N respectively,

with elementary abelian quotient. Therefore, since $P \cap O^p(G)\Phi(P) = P \cap O^p(N)\Phi(P)$, we have, by Tate's result [5, p. 102], that $P \cap O^p(G) = P \cap O^p(N)$, that is

$$G/O^p(G) \cong N/O^p(N).$$

So we can suppose $\Phi(V) = 1$. Consider the chain

$$(3.1) \quad 1 = V_o \triangleleft V_1 \triangleleft \dots \triangleleft V_s = V \trianglelefteq V_{s+1} \trianglelefteq \dots \trianglelefteq V_n = P$$

where $1 = V_o \triangleleft V_1 \triangleleft \dots \triangleleft V_s = V$ is the Φ -chain of V and $V = V_s \trianglelefteq V_{s+1} \trianglelefteq \dots \trianglelefteq V_n = P$ is a chain connecting V and P such that $|V_j/V_{j-1}| = p, j = s + 1, \dots, n$. Since P/V is cyclic, such a chain is uniquely determined. Moreover V_j is weakly closed since V is normal in G . So the chain (3.1) is a strict $p - G$ -chain of P . Then P controls strong fusion by [9] and in particular it controls transfer, that is $G/O^p(G) \cong N/O^p(N)$.

q.e.d.

Corollary 3.3. *Let G be a finite group and p be the smallest prime in $\pi(G)$. Suppose that V is a strongly closed subgroup of $P \in Syl_p(G)$ which possesses a Φ -chain and that P/V is cyclic. Then G is p -nilpotent.*

Proof. By Proposition 3.2 it is enough to show that $N_G(P)$ is p -nilpotent. Let K be a Hall p' -subgroup of $N_G(P)$ which exists by Schur-Zassenhaus Theorem [12, Th. 2.1, p. 221]. K induces automorphisms of V that stabilize the Φ -chain of V since p is the smallest prime in $\pi(G)$. It follows by [12, p. 178] that K induces the identity on $V/\Phi(V)$ and by [12, p. 180] K fixes V . Also K fixes P/V since P/V is cyclic and p is the smallest prime. So K stabilizes the normal chain $1 \trianglelefteq V \trianglelefteq P$ and therefore, again by [12, p. 178], K induces the identity on P . It means that $N_G(P) = P \times K$. q.e.d.

Corollary 3.4. *A finite group G is supersoluble if and only if for all $p \in \pi(G)$, if $P \in Syl_p(G)$, then P possesses a strongly closed subgroup V with a Φ -chain and P/V cyclic.*

Proof. Let G be a minimal counterexample and let r be the smallest prime in $\pi(G)$. Then G has a normal Sylow r -complement K by Corollary 3.3. By minimality of G we have that K is supersoluble and so G has a Sylow tower with respect to the reverse natural order in $\pi(G)$. Let q be the biggest prime in $\pi(G)$ and let $Q \in Syl_q(G)$. Then $Q \trianglelefteq G$. By the hypotheses Q has a strongly closed subgroup V which has Φ -chain and Q/V is cyclic. Since $\Phi(V)$ is strongly closed and subnormal, we have, by Lemma 2.2 (4b) that $\Phi(V) \trianglelefteq G$. Let $\widehat{G} = G/\Phi(V)$ and suppose that $\Phi(V) \neq 1$. Since the hypotheses go to \widehat{G} we have that \widehat{G} is supersoluble and since $\Phi(V) \leq \Phi(G)$ we have that G is supersoluble. Then we can assume $\Phi(V) = 1$. Since Q/V is cyclic with $V \triangleleft G$ and V has a Φ -chain it follows, repeating an argument as in Proposition 3.2, that Q has a strict G -chain $1 = S_o \triangleleft S_1 \triangleleft S_2 \triangleleft \dots \triangleleft S_n = Q$, where $S_i, i = 1, \dots, n$, being weakly closed and subnormal, is normal in G by Lemma 2.2 (4b). Since G/Q is supersoluble by minimality of G we have that G is supersoluble. Vice versa the result is true by Corollary 2 of [8]. q.e.d.

As easy consequence of Corollary 3.4, considering property 3 i) of Lemma 2.2, we have the following:

Corollary 3.5. *A finite group G is supersoluble if and only if there is a normal subgroup N such that G/N has cyclic Sylow subgroups and for every Sylow subgroup P of G we have that $P \cap N$ has a Φ -chain.*

Remarks 3.6.

(1) Now we give two examples which show that if we exclude either that P/V is cyclic or that V has a Φ -chain, then the conclusion in Proposition 3.2 does not hold.

- Consider $G \cong Sym_4$ and let P be a Sylow 2-subgroup of G . Let V be the Klein subgroup which is normal in G . Then P/V is cyclic. On the other hand V does not possess a Φ -chain since an eventually weakly closed subgroup, contained properly in V , would be normal in G by Lemma 2.2 (4b) and this is not the case. On the other hand $N_G(P) = P$. It follows that P does not control transfer, otherwise G would have a normal Sylow 2-complement and this is not true.
- Now consider $G \cong SL(2, 7)$ and let $P \cong Q_{16}$ (the generalized quaternion group) be a Sylow 2-subgroup of G . Take $V = Z(P)$, then V is strongly closed in P and $1 = \Phi(V) \triangleleft V$ is a Φ -chain of V , but P/V is not cyclic. On the other hand $N_G(P) = P$ and so, P does not control transfer, otherwise, as in the previous example, G would have a normal Sylow 2-complement in G and this is not the case.

(2) Now we show that if we suppose that $\Phi(V)$ is weakly closed, instead of strongly closed in the definition of Φ -chain of the strongly closed subgroup V , then the conclusion of Proposition 3.2 is not true.

For, consider $G \cong PSL(2, 17)$ and let $V = P \in Syl_2(G)$. Then P is dihedral of order 16. We have that $\Phi(P)$ is the unique cyclic subgroup of order 4. So it is weakly closed in P but it is not strongly closed. Moreover we observe that there is a unique cyclic subgroup H of order 8 between $\Phi(P)$ and P . Therefore $\Phi(P) \triangleleft H \triangleleft P$ is a chain in which the members are weakly closed and the quotients are of order 2. We have that $N_G(P) = P$ but, as in previous examples, V does not control transfer since, otherwise G would have a normal Sylow 2-complement.

The next lemma extends [10, Th. 19] (see also [15]).

Lemma 3.7. *Let G be a finite group, p the smallest prime in $\pi(G)$ and $P \in Syl_p(G)$. Suppose that any subgroup of P of order p , when p is odd, or any cyclic subgroup of P of order less than or equal to 4, when $p = 2$, is strongly closed in P with respect to G . Then G is p -nilpotent.*

Proof. Let $\Omega(P) = \begin{cases} \Omega_1(P) & \text{for } p \text{ odd} \\ \Omega_2(P) & \text{for } p = 2 \end{cases}$

We have $\Omega_1(P) \leq Z(P)$. Moreover $\Omega_2(P)$, (for $p = 2$), is a product of strongly closed subgroups, then it is strongly closed by Theorem 2 of [1]. If $|\langle x \rangle| = 4$ then $\langle x \rangle \Omega_1(P) \leq Z(P/\Omega_1(P))$ and so $\Omega_2(P) \leq Z_2(P)$. Therefore $\Omega(P)$ has a G -chain, that is $1 \triangleleft \Omega_1(P) = \Omega(P)$ for p odd or $1 \triangleleft \Omega_1(P) \trianglelefteq \Omega_2(P)$ for $p = 2$. We have that $\Omega(P)$ controls strong fusion by [9], therefore $\Omega(P)$ controls transfer, i.e.

$$(3.2) \quad N_G(\Omega(P))/O^p(N_G(\Omega(P))) \cong G/O^p(G)$$

Since any cyclic subgroup of order p , when p is odd, or any cyclic subgroup of order less than or equal to 4, when $p = 2$, is strongly closed in P with respect to $N_G(\Omega(P))$, we have, by induction, that $N_G(\Omega(P))$ is p -nilpotent if $N_G(\Omega(P)) < G$. Then G is p -nilpotent by (3.2).

So we can assume that $\Omega(P) \triangleleft G$. Since all the subgroups of order p , when p is odd, or the cyclic subgroups of order less than or equal to 4, when $p = 2$, are strongly closed and subnormal, then they are normal in G by Lemma 2.2 (4b).

Suppose that $|\langle x \rangle| = p$ with p odd. We have that $|N_G(\langle x \rangle)/C_G(\langle x \rangle)|$ divides $p - 1$. Since p is the smallest prime in $\pi(G)$ we have $C_G(\langle x \rangle) = N_G(\langle x \rangle) = G$. It means $\langle x \rangle \leq Z(G)$. By [13, Satz 5.5, p. 435] we have that G is p -nilpotent.

Now assume that $p = 2$. Clearly if $|\langle x \rangle| = 2$, then $C_G(\langle x \rangle) = G$. So we suppose that $|\langle x \rangle| = 4$. We have $|N_G(\langle x \rangle) : C_G(\langle x \rangle)| \leq 2$. By $N_G(\langle x \rangle) = G$ it follows that $C_G(\langle x \rangle) \trianglelefteq G$. Since all the elements of order less than or equal to 4 of $P \cap C_G(\langle x \rangle)$ are strongly closed in $P \cap C_G(\langle x \rangle)$ with respect $C_G(\langle x \rangle)$, we have, by induction, that $C_G(\langle x \rangle)$ is 2-nilpotent. Suppose that $[G : C_G(\langle x \rangle)] = 2$. Since $O_{2'}(C_G(\langle x \rangle))$ is characteristic in $C_G(\langle x \rangle)$, we have that $O_{2'}(C_G(\langle x \rangle))$ is a normal Sylow 2-complement of G . So G is 2-nilpotent. Therefore we can suppose that all the cyclic subgroups of order less than or equal to 4 are in $Z(G)$. Then, again by [13, Satz 5.5, p. 435] we have that G is 2-nilpotent. q.e.d.

Applying Lemma 3.7 we get a new proof of the following condition for supersolubility. As we have observed in the introduction, this result is a restatement of Theorem 8 in [3].

Corollary 3.8. *Let G be a finite group and suppose that any cyclic subgroup of prime order p (for any $p \in \pi(G)$) or 4 is strongly closed in $P \in Syl_p(G)$ with respect to G . Then G is supersoluble.*

Proof. By Lemma 3.7 G has a Sylow tower with respect to the reverse natural order. Let $P \in Syl_p(G)$ where p is the biggest prime in $\pi(G)$. Then $P \trianglelefteq G$. By induction we have that G/P is supersoluble. Therefore, to prove the result, it is enough to show that each principal factor H/K of G under P has order p . For, let $P_o = \Omega_1(P) = \langle x_i : |x_i| = p \rangle$. Since $\langle x_i \rangle$ is strongly closed and subnormal we have, by Lemma 2.2 (4b) that it is normal in G . Now, with the same proof as [2, Th. 6.7, p. 25] we can prove that $|H/K| = p$. q.e.d.

4. Strict p-G-chains and some applications

A p -subgroup V of a finite group G is said to be normally embedded in G if V is a Sylow p -subgroup of its normal closure (see [4, p. 250]). It is known that if G is a finite p -soluble group and V is a subgroup of $P \in Syl_p(G)$, then V is strongly closed in P with respect to G if and only if it is normally embedded in G (see for instance [11] where this property is stated without a proof for its simplicity). However, for the convenience of the reader, we give an easy proof of it.

Lemma 4.1. *Let G be a finite p -soluble group with $P \in Syl_p(G)$. Then a subgroup V of P is strongly closed in P with respect to G if and only if it is normally embedded in G .*

Proof. First we suppose that V is normally embedded. Then there is a normal subgroup N of G such that $V = P \cap N$ for some $P \in \text{Syl}_p(G)$. So $V^g \cap P = (P \cap N)^g \cap P = P^g \cap N \cap P \leq V$. Therefore V is strongly closed in P with respect to G .

Vice versa, suppose that V is strongly closed in P with respect to G . Then we will show that V is normally embedded in G by induction on $|G|$. If $O_{p'}(G) \neq 1$, then $VO_{p'}(G)/O_{p'}(G)$ is strongly closed in $G/O_{p'}(G)$ by property 3 ii) of Lemma 2.2. So there is $N \trianglelefteq G$ with $N \geq O_{p'}(G)$ such that $VO_{p'}(G)/O_{p'}(G) = (PO_{p'}(G)/O_{p'}(G)) \cap (N/O_{p'}(G)) = (P \cap N)O_{p'}(G)/O_{p'}(G)$. Since $V \leq P \cap N$, it follows that $V = P \cap N$. Therefore we may argue that $O_{p'}(G) = 1$. Since G is p -soluble, it follows that $O_p(G) \neq 1$. By property 3 i) of Lemma 2.2 we have that $V \cap O_p(G)$ is strongly closed in P with respect to G . Moreover, by properties 1) and 4 b) of Lemma 2.2, we have that $V \cap O_p(G) \trianglelefteq G$. Since G is p -soluble and $O_{p'}(G) = 1$, we have, by [12, Th. 3.3, p. 228], that $C_G(O_p(G)) \leq O_p(G)$. So $V \cap O_p(G) \neq 1$. Then, by induction $V/V \cap O_p(G)$ is normally embedded in $G/V \cap O_p(G)$. Therefore V is normally embedded in G . q.e.d.

However the property of Lemma 4.1 is not true in general. Now we give a sufficient condition for a strongly closed subgroup to be normally embedded.

Proposition 4.2. *Let G be a finite group, $p \in \pi(G)$ and $P \in \text{Syl}_p(G)$. Suppose that V is a strongly closed subgroup of P which possesses a strict $p - G$ -chain and $(|N_G(V) : C_G(V)|, p - 1) = 1$. Then $V \in \text{Syl}_p(V^G)$.*

Proof. Let $1 = V_0 \triangleleft V_1 \triangleleft \dots \triangleleft V_n = V$ a strict $p - G$ -chain of V . First observe that the condition $(|N_G(V) : C_G(V)|, p - 1) = 1$ implies $N_G(V) \leq C_G(V_1)$. In fact, suppose that x is a p' -element in $N_G(V)$. Since $V_i, i = 1, \dots, n$, is weakly closed in P , we have that V_i is invariant by x and so it induces an automorphism of $V_i/V_{i-1}, i = 1, \dots, n$. Since $|Aut(V_i/V_{i-1})| = p - 1$ the hypotheses $(|N_G(V) : C_G(V)|, p - 1) = 1$ implies that x centralizes V_i/V_{i-1} for all $i = \{1, \dots, n\}$. Then by [12, p. 178] x centralizes V , in particular $x \in C_G(V_1)$. If x is a p -element, then $x \in Q \in \text{Syl}_p(N_G(V))$. We observe that $Q \in \text{Syl}_p(G)$ and that $V_1 \leq Z(Q)$, so $x \in C_G(V_1)$. If x is any element of $N_G(V)$, then x is a product of a p -element and a p' -element and so we are done.

Now we prove that $V \in \text{Syl}_p(V^G)$ and we suppose that G is a minimal counterexample. We consider $N = N_G(V_1)$ and we distinguish two cases.

- a) $N = G$. If $V_1 = V$ we are done. So we can suppose $V_1 < V$. Then we consider $\bar{G} = G/V_1$ and $\bar{V} = V/V_1$. Since $N_{\bar{G}}\bar{V} = N_G(V)/V_1$ by Lemma 2.2 and considering that $C_{\bar{G}}(\bar{V}) \geq C_G(V)/V_1$ we have that \bar{G} satisfies the same hypotheses as G . Therefore \bar{V} is a Sylow p -subgroup of $(\bar{V})^{\bar{G}}$. But $(\bar{V})^{\bar{G}} = V^G/V_1$ and then V is a Sylow p -subgroup of V^G .
- b) $N < G$. Obviously $V \leq P \leq N$, so the hypotheses go to N . It follows that $V \in \text{Syl}_p(V^N)$. Now let $g \in N \setminus C_G(V_1)$. By Theorem A of [7] we have that V^N has a normal Sylow p -complement K . Since K is characteristic in V^N we have that K is normal in N . So $K \leq C_G(V_1)$. Since $V^g \in \text{Syl}_p(V^N)$ we have $V^g = V^k$ where $k \in K$. It follows $gk^{-1} \in N_G(V)$. Then $g = nk$ with $n \in N_G(V) \leq C_G(V_1)$. So $g \in C_G(V_1)$ against the assumption. It follows $N_G(V_1) = C_G(V_1)$.

This condition implies that if $\langle x \rangle = V_1$ then x is not conjugate to its powers. Furthermore, since V_1 is weakly closed in P with respect to G , we have that x does not commute with its conjugates. So either by [6, Proposition 2.2] for the case p odd or by the Z^* -Theorem of Glauberman if $p = 2$ (see [16, p. 315]) we have that $V_1 O_{p'}(G)/O_{p'}(G) \leq Z(G/O_{p'}(G))$. Observe that, since V_1 is not normal in G , we can suppose $O_{p'}(G) \neq 1$. So, by minimality of G we get that $VO_{p'}(G)/O_{p'}(G)$ is a Sylow p -subgroup of $(VO_{p'}(G)/O_{p'}(G))^{G/O_{p'}(G)}$. Let $M/O_{p'}(G) = (VO_{p'}(G)/O_{p'}(G))^{G/O_{p'}(G)}$. Since $M \geq V^G$ and $(|V|, |O_{p'}(G)|) = 1$, we have that $V \in \text{Syl}_p(V^G)$ and so we are done.

q.e.d.

Corollary 4.3. *Let G be a finite group and $P \in \text{Syl}_p(G)$. Suppose that $(|G|, p - 1) = 1$ and that V is strongly closed in P with a strict $p - G$ -chain. Then $VO_{p'}(G)$ is normal in G .*

Proof. By Proposition 4.2 we have that V is a Sylow p -subgroup of V^G . We have that $N_G(V)/C_G(V)$ is a p -group. In fact suppose that x is a p' -element of $N_G(V)$. If $1 = V_0 \trianglelefteq \dots \trianglelefteq V_n = V$ is a $p - G$ -chain of V , we have that V_i is invariant by x since V_i is weakly closed in P . So x induces an automorphism of V_{i+1}/V_i , $(i = 0, \dots, n - 1)$, of order dividing $p - 1$. Then, by assumption x induces the identity. It follows, by [12, p. 178] that x induces the identity on V , that is $x \in C_G(V)$. Then by [7] we have that V^G has a normal Sylow p -complement, say K . Of course $K \leq O_{p'}(G)$. If $O_{p'}(G) = 1$, then $V = V^G$ and we are done. So we may assume that $O_{p'}(G) \neq 1$. But $G/O_{p'}(G)$ satisfies the same hypotheses as G . So by induction $(VO_{p'}(G)/O_{p'}(G))O_{p'}(G/O_{p'}(G)) = VO_{p'}(G)/O_{p'}(G)$ is normal in $G/O_{p'}(G)$, that is $VO_{p'}(G) \trianglelefteq G$.

q.e.d.

Remarks 4.4. *We prove that if we exclude one of the two hypotheses in Proposition 4.2, then the conclusion is not true.*

- a) Consider the simple group $PSL(2, 17)$. Then the Sylow p -subgroups for p odd are cyclic. In particular a Sylow 3-subgroup P has order 9. Let V be a subgroup of order 3. Then V is strongly closed and obviously has a $p - G$ -chain. However V is not a Sylow 3-subgroup of its normal closure. The reason depends on the fact that $|N_G(V) : C_G(V)|, 3 - 1 \neq 1$.
- b) Let $G \cong Sz(2^{2n+1})$ be a Suzuki group and $P \in \text{Syl}_2(G)$. Then $Z(P) = \Omega_1(P)$ and so $V = Z(P)$ is strongly closed in P with respect to G . Here, obviously we have $(|N_G(V) : C_G(V)|, 2 - 1) = 1$, but V is not a Sylow 2-subgroup of V^G . The reason depends on the fact that V has not strict $p - G$ -chain although it has obviously a $p - G$ -chain.

REFERENCES

[1] M. Aschbacher, The generalized Fitting subsystem of a fusion system, *Mem. Amer. Math. Soc.*, **209** no. 986 (2011) vi+110.

- [2] H. G. Bray, W. E. Deskins, D. Johnson, J. F. Humphreys, B. M. Puttaswamaiah, P. Venzke and G. L. Walls, *Between Nilpotent and Solvable*, Edited and with a preface by Michael Weinstein, Polygonal Publ. House, Washington, N. J., 1982.
- [3] P. Csörgö and M. Herzog, On supersolvable groups and the nilpotator, *Comm. Algebra*, **32** no. 2 (2004) 609–620.
- [4] K. Doerk and T. Hawkes, *Finite Soluble Groups*, de Gruyter Expositions in Mathematics, **4**, Walter de Gruyter & Co., Berlin, 1992.
- [5] L. Dornhoff, *Group Representation Theory, Part A: Ordinary Representation Theory*, Pure and Applied Mathematics, **7**, Marcel Dekker Inc., New York, 1971.
- [6] R. J. Flores and R. M. Foote, Strongly closed subgroups of finite groups, *Adv. Math.*, **222** no. 2 (2009) 453–484.
- [7] A. L. Gilotti and L. Serena, On the existence of normal Sylow p -complement, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)*, **69** no. 5 (1980) 228–231.
- [8] A. L. Gilotti and L. Serena, Sottogruppi massimali dei sottogruppi di Sylow e complementi normali, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)*, **76** no. 3 (1984) 161–166.
- [9] A. L. Gilotti and L. Serena, Strongly closed subgroups and strong fusion, *J. London Math. Soc. (2)*, **32** no. 1 (1985) 103–106.
- [10] A. L. Gilotti and L. Serena, Remarks on chains of weakly closed subgroups, *J. Group Theory*, **14** no. 6 (2011) 807–818.
- [11] D. M. Goldschmidt, Strongly closed 2-subgroups of finite groups, *Ann. of Math. (2)*, **102** no. 3 (1975) 475–489.
- [12] D. Gorenstein, *Finite groups*, Harper & Row Publishers, New York, 1968.
- [13] B. Huppert, *Endliche Gruppen I*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **134**, Springer-Verlag, Berlin, 1967.
- [14] B. Huppert and N. Blackburn, *Finite Groups III*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **243**, Springer-Verlag, Berlin, 1982.
- [15] S. R. Li, On minimal subgroups of finite groups, *Comm. Algebra*, **22** no. 6 (1994) 1913–1918.
- [16] M. Suzuki, *Groups Theory II*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **248**, Springer-Verlag, New York, 1986.

Anna Luisa Gilotti

Dipartimento di Matematica, Università di Bologna, p.zza di Porta S. Donato 5, 40126 Bologna, Italy

Email: gilotti.isa@gmail.com

Luigi Serena

Dipartimento di Matematica e Informatica U.Dini, Università di Firenze, Viale Morgagni 67/A, 50134 Firenze, Italy

Email: luigi.serena@unifi.it