



<http://ijgt.ui.ac.ir>

International Journal of Group Theory
 ISSN (print): 2251-7650, ISSN (on-line): 2251-7669
 Vol. x No. x (202x), pp. xx-xx.
 © 202x University of Isfahan



www.ui.ac.ir

THE INFLUENCE OF \mathcal{H} -SUBGROUPS ON p -NILPOTENCY AND p -SUPERSOLVABILITY OF FINITE GROUPS

QUANFU YAN* AND ZHENCAI SHEN

ABSTRACT. Let G be a finite group. A subgroup H of G is an \mathcal{H} -subgroup in G if $N_G(H) \cap H^g \leq H$ for any $g \in G$. In this article, by using the concept of \mathcal{H} -subgroups, we study the influence of the intersection of $O^p(G_p^*)$ and the members of some fixed $\mathcal{M}_d(P)$ on the structure of the group G , where P is a Sylow p -subgroup of G . Some new criteria for a group to be p -nilpotent and p -supersolvable are given and some recent results are extended and generalized.

1. Introduction

Throughout this article, all groups are finite. Let G be a group and p a prime. Let $O^p(G)$ be the unique smallest normal subgroup of G for which the corresponding factor group is a p -group. Following Berkovich and Isaacs in [2], we let G_p^* be the unique smallest normal subgroup of G for which the corresponding factor group is abelian of exponent dividing $p - 1$. A group G is called p -supersolvable if every chief factor of G either has order p or it has order coprime to p . It is known that G is p -supersolvable if and only if G_p^* is p -nilpotent (see [2, Lemma 3.6]). To state our results, we need to recall some concepts.

Goldschmidt, Flores and Foote in [5, 6, 7, 8] studied the following concept.

Definition 1.1. *Let $K \leq G$. A subgroup H of K is called strongly closed in K with respect to G if for every element h of H , $h^G \cap K \subseteq H$, where h^G denotes the G -conjugacy class of h . In particular, when*

Communicated by Hamid Mousavi.

MSC(2010): Primary: 20D20; Secondary: 20D25.

Keywords: \mathcal{H} -subgroup, weakly \mathcal{H} -subgroup, p -supersolvability, p -nilpotency.

Article Type: Research Paper.

Received: 21 September 2022, Accepted: 01 January 2023.

*Corresponding author.

H is a subgroup of prime-power order and K is a Sylow subgroup containing it, H is simply said to be a strongly closed subgroup.

This concept plays an important role in group theory and representation theory, particularly in the case when K is a Sylow subgroup of G . If H is a p -subgroup, then H is strongly closed in a Sylow p -subgroup if and only if it is strongly closed in $N_G(P)$, so this concept for a p -subgroup does not depend on the Sylow subgroup containing it. In 2000, Bianchi et al. in [3] introduced the definition of \mathcal{H} -subgroups which is related to strongly closed subgroups. A subgroup H of a group G is an \mathcal{H} -subgroup in G if $N_G(H) \cap H^g \leq H$ for any $g \in G$. Notice that if H is a subgroup of G of prime-power order, then H is an \mathcal{H} -subgroup in G if and only if H is a strongly closed subgroup. Asaad et al. in [1] introduced the following subgroup embedding property called weakly \mathcal{H} -subgroup, which is a generalization of \mathcal{H} -subgroup. A subgroup H of a group G is called a weakly \mathcal{H} -subgroup in G if there exists a normal subgroup N of G such that $G = HN$ and $H \cap N \in \mathcal{H}(G)$, where $\mathcal{H}(G)$ denotes the set of all \mathcal{H} -subgroups in G . In addition, they proved the following result.

Theorem 1.2. [1] *Let p be the smallest prime dividing the order of G and let P be a Sylow p -subgroup of G . If all maximal subgroups of P are weakly \mathcal{H} -subgroups in G , then G is p -nilpotent.*

In general, a weakly \mathcal{H} -subgroup in G is not always an \mathcal{H} -subgroup. However, in Lemma 2.5, we can show that if H is a p -subgroup of G , then $H \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G if and only if $H \cap O^p(G_p^*)$ is a weakly \mathcal{H} -subgroup in G . In Lemma 2.6, we prove that if H is a maximal subgroup of a Sylow p -subgroup of G , then $H \cap O^p(G)$ is an \mathcal{H} -subgroup in G if and only if H is a weakly \mathcal{H} -subgroup in G .

Let P be a nontrivial finite p -group and $\mathcal{M}(P)$ be the set of all maximal subgroups of P . Li et al. in [12] introduced a subset of $\mathcal{M}(P)$ as follows: Let d be the smallest generator number of a p -group P and $\mathcal{M}_d(P) = \{P_1, P_2, \dots, P_d\}$ be a set of maximal subgroups of P such that $\Phi(P) = \bigcap_{i=1}^d P_i$. For a fixed P , $\mathcal{M}_d(P)$ is not unique in general. We know that $|\mathcal{M}(P)| = \frac{p^d - 1}{p - 1}$ and $|\mathcal{M}_d(P)| = d$. Hence, it is easy to see that $\mathcal{M}_d(P)$ is a very small subset of $\mathcal{M}(P)$ while d is large enough. In [13], Shen et al. obtained the following criteria of p -nilpotency by using the set $\mathcal{M}_d(P)$ and the concept of \mathcal{H} -subgroups.

Theorem 1.3. [13] *Let p be a prime dividing the order of a group G , and P a Sylow p -subgroup of G . Then G is p -nilpotent if and only if $N_G(P)$ is p -nilpotent and every member of some fixed $\mathcal{M}_d(P)$ is an \mathcal{H} -subgroup of G .*

Theorem 1.4. [13] *Let p be a prime dividing the order of a group G with $(|G|, p - 1) = 1$ and let P be a Sylow p -subgroup of G . Then the following statements are equivalent:*

- (1) G is p -nilpotent.
- (2) For every member of $\mathcal{M}(P)$ is an \mathcal{H} -subgroup of G .
- (3) For every member of some fixed $\mathcal{M}_d(P)$ is an \mathcal{H} -subgroup of G .

Our goal in this article is to improve and extend the results we mentioned above. Our basic idea is from several recent articles (for example, see [10, 15, 16, 17]) where p -supersolvability or p -nilpotency of a group G is obtained by assuming that some p -subgroups of $O^p(G)$ or $O^p(G_p^*)$ satisfy some embedding property in G . By using the concept of \mathcal{H} -subgroups, we investigate the influence of the intersection of $O^p(G_p^*)$ and the members of some fixed $\mathcal{M}_d(P)$ on the structure of the group G , where P is a Sylow p -subgroup of G . We obtain the following theorems.

Theorem A. *Let G be a p -solvable group and P be a Sylow p -subgroup of G , where p is a prime divisor of the order of G . Then G is p -supersolvable if and only if for every member P_i of some fixed $\mathcal{M}_d(P)$, $P_i \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G .*

Theorem B. *Let p be a prime dividing the order of a group G and P a Sylow p -subgroup of G . Then the following statements are equivalent:*

- (1) G is p -nilpotent.
- (2) $N_G(P)$ is p -nilpotent and for every member P_i of some fixed $\mathcal{M}_d(P)$, P_i is a weakly \mathcal{H} -subgroup in G .
- (3) $N_G(P)$ is p -nilpotent and for every member P_i of some fixed $\mathcal{M}_d(P)$, $P_i \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G .

Theorem C. *Let p be a prime dividing the order of a group G with $(|G|, p - 1) = 1$ and let P be a Sylow p -subgroup of G . Then the following statements are equivalent:*

- (1) G is p -nilpotent.
- (2) For every member P_i of some fixed $\mathcal{M}_d(P)$, P_i is a weakly \mathcal{H} -subgroup of G .
- (3) For every member P_i of some fixed $\mathcal{M}_d(P)$, $P_i \cap O^p(G_p^*) = P_i \cap O^p(G)$ is an \mathcal{H} -subgroup in G .

2. Preliminaries

Lemma 2.1. *Let G be a group and let H, K, N be subgroups of G satisfying $H \in \mathcal{H}(G)$, $H \leq K$, and N is normal in G . Then the following hold:*

- (1) $H \in \mathcal{H}(K)$;
- (2) If P is a Sylow p -subgroup of G , then $P \cap N \in \mathcal{H}(G)$;
- (3) If H is subnormal in K , then H is normal in K ;
- (4) If $N \leq H$, then $H \in \mathcal{H}(G)$ if and only if $H/N \in \mathcal{H}(G/N)$;
- (5) If H is a p -subgroup of G such that $(|H|, |N|) = 1$, then $HN \in \mathcal{H}(G)$ and $HN/N \in \mathcal{H}(G/N)$.
- (6) If P is a Sylow p -subgroup of N , then $P \in \mathcal{H}(G)$.

Proof. For (1)-(5), see [1, Lemma 2.1]. For (6), see [3, Proposition 3(2)]. □

Lemma 2.2. [15, Lemma 2.9][16, Lemma 2.4] *Let p be a prime dividing the order of a finite group G , $H \leq G$ and $N \trianglelefteq G$. Then $O^p(H_p^*) \leq O^p(G_p^*)$, $O^p(H) \leq O^p(G)$, $(G/N)_p^* = G_p^*N/N$, $O^p(G/N) = O^p(G)N/N$ and $O^p((G/N)_p^*) = O^p(G_p^*)N/N$.*

Lemma 2.3. *Let P be a Sylow p -subgroup of G for a prime p . If H and K are strongly closed in P with respect to G , then $H \cap K$ is strongly closed in P with respect to G .*

Proof. Let $a \in H \cap K$. By assumption, $a^G \cap P \subseteq H$ and $a^G \cap P \subseteq K$. Hence, $a^G \cap P \subseteq H \cap K$. It follows that $H \cap K$ is strongly closed in P with respect to G . \square

Lemma 2.4. *Let P be a Sylow p -subgroup of G for a prime p and let N be a normal subgroup of G . If $H \leq P$ is an \mathcal{H} -subgroup in G , then $H \cap N \in \mathcal{H}(G)$.*

Proof. Notice that if H is a p -subgroup of G , then H is an \mathcal{H} -subgroup in G if and only if H is a strongly closed subgroup. By Lemma 2.1(2), $P \cap N \in \mathcal{H}(G)$ and so $P \cap N$ is strongly closed subgroup. By Lemma 2.3, $H \cap N = (P \cap N) \cap H$ is strongly closed subgroup and hence $H \cap N \in \mathcal{H}(G)$, as wanted. \square

By the above lemma, if a p -subgroup H of G is an \mathcal{H} -subgroup in G , then $H \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G . But the converse is not true. Consider $G = S_4$ and its 2-subgroup $H = \langle (13) \rangle$ for instance. It is clear that $G_2^* = G$ and so $O^2(G_2^*) = A_4$. Hence, $H \cap O^2(G_2^*) = 1$ is an \mathcal{H} -subgroup in G . Let P be a Sylow 2-subgroup of G containing H . If $H \in \mathcal{H}(G)$, then by Lemma 2.1(1)(3), H is normal in P . This is a contradiction. Hence, H is not an \mathcal{H} -subgroup in G . However, we have the following:

Lemma 2.5. *Let p be a prime dividing the order of a finite group G and H be a p -subgroup of G . Then $H \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G if and only if $H \cap O^p(G_p^*)$ is a weakly \mathcal{H} -subgroup in G .*

Proof. We only need to prove the sufficiency. Assume that $H \cap O^p(G_p^*)$ is a weakly \mathcal{H} -subgroup in G . Then by definition, there is a normal subgroup N of G such that $G = (H \cap O^p(G_p^*))N$ and $H \cap O^p(G_p^*) \cap N$ is an \mathcal{H} -subgroup in G . Notice that $G/N \cong H \cap O^p(G_p^*) / H \cap O^p(G_p^*) \cap N$ is a p -group. It follows that $O^p(G) \leq N$. By Lemma 2.2, we have $O^p(G_p^*) \leq O^p(G)$. Since $O^p(G_p^*) \trianglelefteq G$, by Lemma 2.4, $H \cap O^p(G_p^*) = H \cap O^p(G_p^*) \cap N$ is an \mathcal{H} -subgroup in G , as wanted. \square

Lemma 2.6. *Let p be a prime dividing the order of a group G and P a Sylow p -subgroup of G . If H is a maximal subgroup of P , then $H \cap O^p(G)$ is an \mathcal{H} -subgroup in G if and only if H is a weakly \mathcal{H} -subgroup in G .*

Proof. If H is a weakly \mathcal{H} -subgroup in G , then there exists a normal subgroup N of G such that $G = HN$ and $H \cap N$ is an \mathcal{H} -subgroup in G . Notice that $O^p(G) \leq N$. Then by Lemma 2.4, $H \cap O^p(G)$ is an \mathcal{H} -subgroup in G . Now assume that $H \cap O^p(G)$ is an \mathcal{H} -subgroup of G . Consider $M = HO^p(G)$. If $G = M$, then by definition, H is a weakly \mathcal{H} -subgroup in G . If $M < G$, then it is clear that $|G : M| = p$ and H is a Sylow p -subgroup of M . Notice that $M \trianglelefteq G$. Hence, H is an \mathcal{H} -subgroup in G by Lemma 2.1(6). Clearly, H is a weakly \mathcal{H} -subgroup in G . \square

Lemma 2.7. [15, Lemma 2.8] *Let p be a prime dividing the order of a finite group G and H be a p -subgroup of G . Let $L \trianglelefteq G$ and N be a normal p' -subgroup of G . Then $HN/N \cap LN/N = (H \cap L)N/N$.*

3. Main results

Proof of Theorem A. Suppose that G is p -supersolvable. Then G_p^* is p -nilpotent and so G_p^* has a normal subgroup K such that $G_p^* = P \rtimes K$ and $P \cap K = 1$. It is clear that $K = O^p(G_p^*)$. Hence, for any $P_i \in \mathcal{M}_d(P)$, $P_i \cap O^p(G_p^*) = 1$ is an \mathcal{H} -subgroup in G .

Conversely, suppose that the theorem is not true and let G be a counterexample with minimal order. Set $\mathcal{M}_d(P) = \{P_1, P_2, \dots, P_d\}$. We will obtain a contradiction by the following steps.

Step 1. $O_{p'}(G) = 1$.

Assume $O_{p'}(G) > 1$ and consider $G/O_{p'}(G)$. By Lemma 2.1(5), Lemma 2.2 and Lemma 2.7, $G/O_{p'}(G)$ satisfies the hypothesis of this theorem. By the minimal choice of G , we have that $G/O_{p'}(G)$ is p -supersolvable, and thus G is p -supersolvable, which is a contradiction. Hence, $O_{p'}(G) = 1$.

Step 2. Let $M = O^p(G_p^*)$ and $N = O_p(M)$. Then $N > 1$ and $C_M(N) \leq N$.

Since G is not p -supersolvable, it follows that G_p^* is not p -nilpotent. If $M = 1$, then G_p^* is a p -group, a contradiction. Hence, $M > 1$. Since $1 < M \leq G$ is p -solvable and $O_{p'}(M) \leq O_{p'}(G) = 1$, we have $N > 1$ and $C_M(N) \leq N$ by Hall-Higman's Lemma (see [11, Theorem 3.21]).

Step 3. $G_p^* \leq C_G(N)$.

Notice that for each $i \in \{1, \dots, d\}$, $P_i \cap O^p(G_p^*)$ is an \mathcal{H} -subgroup in G . Since $N \leq O^p(G_p^*)$, by Lemma 2.4, for each i , $P_i \cap N = (P_i \cap O^p(G_p^*)) \cap N \in \mathcal{H}(G)$. Since $P_i \cap N \trianglelefteq N \trianglelefteq G$, by Lemma 2.1(3), we have that $P_i \cap N \trianglelefteq G$. Hence, $\Phi(P) \cap N = \bigcap_{i=1}^d (P_i \cap N) \trianglelefteq G$.

Let $L = \Phi(P) \cap N$. If $L > 1$, then it is clear that G/L satisfies the hypothesis of this theorem. Hence, G/L is p -supersolvable. As $L \trianglelefteq G$ and $L \leq \Phi(P)$, we have $L \leq \Phi(G)$. Hence, $G/\Phi(G)$ is p -supersolvable. Since the class of p -supersolvable groups is a saturated formation, it follows that G is p -supersolvable, a contradiction. Thus, $L = 1$.

If $N \leq P_i$, then $[N, G_p^*] \leq N = P_i \cap N$. Assume now $N \not\leq P_i$. Then $P = P_i N$ and so $|P : P_i| = |N : P_i \cap N| = p$. Consider $\overline{G} = G/P_i \cap N$. Notice that $\overline{G}/C_{\overline{G}}(\overline{N})$ is isomorphic to a subgroup of $\text{Aut}(\overline{N}) \cong C_{p-1}$. It follows that $(\overline{G})_p^* \leq C_{\overline{G}}(\overline{N})$. By Lemma 2.2, we have $(\overline{G})_p^* = \overline{G}_p^*$ and so $[G_p^*, N] \leq P_i \cap N$. Hence, $[N, G_p^*] \leq P_i \cap N$ for each i . It follows that $[N, G_p^*] \leq \Phi(P) \cap N = 1$ and so $G_p^* \leq C_G(N)$.

Step 4. Final contradiction.

By Step 2, $C_M(N) \leq N$. Notice that $M \leq G_p^*$. Then by Step 3, $M \leq C_M(N)$ and hence $N = M$. Since $O^p(G_p^*)/N = O^p(G_p^*)/M \cong O^p(G_p^*)O^p(G)/O^p(G) \leq G/O^p(G)$ and N are p -groups, it follows that $O^p(G_p^*)$ is a p -group and so is G_p^* , a contradiction. \square

Proof of Theorem B. Clearly, we only need to show that (3) implies (1). We always set $\mathcal{M}_d(P) = \{P_1, P_2, \dots, P_d\}$. Suppose that it is not true and let G be a counterexample with minimal order. We work in the following steps to obtain a contradiction.

Step 1. $O_{p'}(G) = 1$.

Assume that $O_{p'}(G) > 1$. Consider $\bar{G} = G/O_{p'}(G)$. Notice that $N_{\bar{G}}(\bar{P}) = \overline{N_G(P)}$ is p -nilpotent. Then by Lemma 2.1(5), Lemma 2.2 and Lemma 2.7, $G/O_{p'}(G)$ satisfies the hypotheses of this theorem. By the minimal choice of G , we have that $G/O_{p'}(G)$ is p -nilpotent, and thus G is p -nilpotent, which is a contradiction. Hence, $O_{p'}(G) = 1$.

Step 2. If $P \leq H < G$, then H is p -nilpotent.

Noting that $N_H(P) \leq N_G(P)$, we have $N_H(P)$ is p -nilpotent. By Lemma 2.1(1), $P_i \cap O^p(G_p^*) \in \mathcal{H}(H)$ for each i . It is clear that $O^p(H_p^*) \trianglelefteq H$ and $O^p(H_p^*) \leq O^p(G_p^*)$ by Lemma 2.2. Thus, $P_i \cap O^p(H_p^*) \in \mathcal{H}(H)$ by Lemma 2.4. Hence, by induction, H is p -nilpotent.

Step 3. G is p -solvable.

Since G is not p -nilpotent, by a result of Thompson [14, Corollary], there exists a non-trivial characteristic subgroup T of P such that $N_G(T)$ is not p -nilpotent. Choose T such that the order of T is as large as possible. Since $N_G(P)$ is p -nilpotent, $N_G(K)$ is p -nilpotent for any characteristic subgroup K of P satisfying $T < K \leq P$. Notice that $T \text{ char } P \trianglelefteq N_G(P)$. Then $T \trianglelefteq N_G(P)$ and so $N_G(P) \leq N_G(T)$. By Step 2, $N_G(T) = G$ and hence $T \leq O_p(G)$. By the choice of T , we have $T = O_p(G)$. Now applying the result of Thompson again, we have that $G/O_p(G)$ is p -nilpotent, therefore G is p -solvable.

Step 4. Final contradiction.

By Theorem A, G is p -supersolvable. In particular, G is p -solvable with p -length 1. By Step 1, we have $P \trianglelefteq G$ and so $G = N_G(P)$ is p -nilpotent, a contradiction. \square

Proof of Theorem C. We only need to show that (3) implies (1). Since $(|G|, p-1) = 1$, we have that $G_p^* = G$ and $O^p(G_p^*) = O^p(G)$. It is easy to see that the hypotheses are inherited by $N_G(P)$. If $N_G(P) < G$, then $N_G(P)$ is p -nilpotent by induction. It follows from Theorem B that G is p -nilpotent, as wanted. Assume that $N_G(P) = G$. Then $P \trianglelefteq G$ and so G is p -solvable. By Theorem A, we have that G is p -supersolvable. Hence, $G = G_p^*$ is p -nilpotent, as wanted. \square

4. Applications

Recently, Chen et al in [4] proved the following theorem.

Theorem 4.1. [4] Let G be a group and let P be a Sylow p -subgroup of G , where p is the smallest prime dividing the order of G . If all maximal subgroups of P are weakly \mathcal{H} -subgroups in $N_G(P)$ and P' is normal in G , then G is p -nilpotent.

In view of Theorem A, Theorem B and Theorem C, we extend the above theorem and obtain the following result.

Corollary 4.2. Let p be a prime dividing the order of a group G with $(|G|, p-1) = 1$ and $P \in \text{Syl}_p(G)$. If for every member P_i of some fixed $\mathcal{M}_d(P)$, $P_i \cap O^p(N_G(P))$ is an \mathcal{H} -subgroup in $N_G(P)$ and P' is normal in G , then G is p -nilpotent.

Proof. By Theorem C, we have that $N_G(P)$ is p -nilpotent and so $N_G(P) = P \times O_{p'}(N_G(P)) = P \times O^p(N_G(P))$. Hence, $P_i \cap O^p(N_G(P)) = 1$. If $P' = 1$, then $C_G(P) = N_G(P)$. By Burnside's theorem ([9, Theorem 7.4.3]), G is p -nilpotent. Now assume $P' > 1$. Consider $\bar{G} = G/P'$. Clearly, $P' \leq \Phi(P)$ and $O^p(N_{\bar{G}}(\bar{P}))$ is a p' -group. Then $\bar{P}_i \cap O^p(N_{\bar{G}}(\bar{P})) = 1$ is an \mathcal{H} -subgroup of $N_{\bar{G}}(\bar{P})$. It follows that \bar{G}' satisfies the hypotheses. By induction, G/P' is p -nilpotent. Notice that $P' \leq \Phi(P)$ and $P' \trianglelefteq G$. Then $P' \leq \Phi(G)$ and so $G/\Phi(G)$ is p -nilpotent. It follows that G is p -nilpotent. \square

Corollary 4.3. Let p be a prime dividing the order of a group G with $(|G|, p-1) = 1$ and $P \in \text{Syl}_p(G)$. If for every member P_i of some fixed $\mathcal{M}_d(P)$, P_i is a weakly \mathcal{H} -subgroup in $N_G(P)$ and P' is normal in G , then G is p -nilpotent.

Acknowledgments

The authors wish to thank the referee for careful reading of the article and detailed report including corrections and comments.

REFERENCES

- [1] M. Asaad, A. A. Heliel and M. M. Al-Mosa Al-Shomrani, On weakly \mathcal{H} -subgroups of finite groups, *Comm. Algebra*, **40** (2012) 3540–3550.
- [2] Y. Berkovich and I. M. Isaacs, p -Supersolvability and actions on p -groups stabilizing certain subgroups, *J. Algebra*, **414** (2014) 82–94.
- [3] M. Bianchi, A. Gillio Berta Mauri, M. Herzog and L. Verardi, On finite solvable groups in which normality is a transitive relations, *J. Group Theory*, **3** (2000) 147–156.
- [4] R. Chen, X. Li and X. Zhao, On weakly \mathcal{H} -subgroups of finite groups II*, *Comm. Algebra*, **50** (2022) 4009–4015.
- [5] R. J. Flores and R. M. Foote, Strongly closed subgroups of finite groups, *Adv. Math*, **222** (2009) 453–484.
- [6] D. Goldschmidt, 2-fusion in finite groups, *Ann. of Math*, **99** (1974) 70–117.
- [7] D. Goldschmidt, Strongly closed 2-subgroups of finite groups, *Proc. Conf. on Finite Groups (Univ. Utah, Park City, Utah)*, (1975) 109–110.
- [8] D. Goldschmidt, Strongly closed 2-subgroups of finite groups, *Ann. of Math*, **103** (1975) 475–489.

- [9] D. Gorenstein, *Finite Groups*, Harper and Row, New York-London, 1968
- [10] Y. Guo and I. M. Isaacs, Conditions on p -subgroups implying p -nilpotence or p -supersolvability, *Arch. Math*, **105** (2015) 215–222.
- [11] I. M. Isaacs, *Finite Group Theory*, **92**, American Mathematical Society, Providence, 2008
- [12] S. Li and X. He, On normally embedded subgroups of prime power order in finite groups, *Comm. Algebra*, **36** (2008) 2333–2340.
- [13] Z. Shen and N. Du, Finite Groups with \mathcal{H} -Subgroups, *Algebra Colloq.*, **20** (2013) 421–426.
- [14] J. G. Thompson, Normal p -complements for finite groups, *J. Algebra*, **1** (1964) 43–46.
- [15] H. Yu, Some sufficient and necessary conditions for p -supersolvability and p -nilpotence of a finite group, *J. Algebra Appl*, **16** (2017) 1750052.
- [16] H. Yu, X. Xu and G. Zhang, On generalized $S\Phi$ -supplemented subgroups of finite groups, *J. Algebra Appl*, **18** (2019) 1950204.
- [17] H. Yu, X. Xu and G. Zhang, A note on S -semipermutable and S -permutably embedded subgroups of finite groups, *Ricerche mat.*, **105** (2022). <https://doi.org/10.1007/s11587-022-00717-1>.

Quanfu Yan

Department of Mathematical Sciences, Kent State University, Kent, USA

Email: qyan5@kent.edu

Zhencai Shen

Department of Mathematics, China Agricultural University, Beijing, China

Email: zhencai688@sina.com